

Classification of Singular Fibres on Rational Elliptic Surfaces in Characteristic Three

Tyler Jarvis, William E. Lang,
Nansen Petrosyan, Gretchen Rimmasch,
Julie Rogers, Erin D. Summers

February 1, 2008

1 Introduction

1.1 Background and Overview

In [7] U. Persson classified all possible configurations of singular fibres on rational elliptic surfaces over the complex numbers. He found 279 different configurations. R. Miranda [4] used a more combinatorial method to redo Persson's list. W. Lang [2] continued the classification for characteristic two. Following the method of Lang, we list all possible configurations of singular fibres on rational elliptic surfaces in characteristic three. In total, we find that 267 distinct configurations exist.

Let k be an algebraically closed field of characteristic three. Let $f : X \rightarrow \mathbb{P}^1$ be a rational elliptic surface over k . We will assume that the surface is relatively minimal and that f has a section. Such a surface has an expression in terms of a Weierstrass equation

$$y^2 = x^3 + b_2x^2 - b_4x + b_6, \quad (1.1.1)$$

with $b_2, b_4, b_6 \in k[t]$ and $\deg b_i \leq i$. The discriminant Δ for this surface is

$$\Delta = -b_2^2(b_2b_6 - b_4^2) + b_4^3. \quad (1.1.2)$$

Assuming that the surface has good reduction at infinity, the discriminant Δ of the surface is a polynomial in t of degree exactly 12. According to Tate's algorithm [9], the surface has a singular fibre of multiplicative type at a given point $t = p$ of the base if and only if $\Delta(p) = 0$ but $b_2(p) \neq 0$, and a singular fibre of additive type at $t = p$ if and only if $\Delta(p) = 0$ and $b_2(p) = 0$. There are four general configurations of singular fibres that we will consider:

1. $b_2 \neq 0$ and all the singular fibres are multiplicative.
2. $b_2 \neq 0$ and one of the singularities is additive, and any remaining singularities are multiplicative.

3. $b_2 \neq 0$ and two of the singularities are additive, and any remaining singularities are multiplicative.
4. $b_2 \equiv 0$, in which case all the singularities are additive, and there can be more than two distinct additive singularities. In this case the j -invariant is also identically zero.

We will treat the first case separately and the other three together. The structure of the paper will be as follows. First, some preliminaries relating to all surfaces will be discussed. We will then consider those surfaces which have only multiplicative type singular fibres, followed by those surfaces with at least one additive type singular fibre. Finally we will give a summary of the results.

1.2 Acknowledgments

We are grateful to Heidi Jarvis for her expert help with proofreading and typesetting. Research of the first author was supported in part by NSF grant DMS-0105788. The last four authors were supported in part by graduate research assistantships from the Brigham Young University mathematics department.

2 Preliminaries

Let $f : X \rightarrow \mathbb{P}^1$ be a rational elliptic surface over an algebraically closed field k of characteristic three. We assign several invariants to each singular fibre, as given in Table 1. First we list the Kodaira type F . But, unlike in characteristic zero, each singularity may appear in different ways, for example with different orders of vanishing of Δ , so we also list Lang's case number L from the classification in [3] of isolated additive type singular fibres on a rational elliptic surface in characteristic three. Lang's normal forms from that classification play a key role in this project.

The order of vanishing of Δ is denoted by δ . Associated with each singular fibre is a lattice Λ of rank r , generated by the irreducible components of the singular fibre which do not meet the zero section. The lattice Λ is a sublattice of the Nèron-Severi lattice of the entire surface, with discriminant d , and is orthogonal to the lattices associate to the other fibres.

Table 1: Fibre Invariants

F	L	δ	r	Λ	d
I_n		n	$n-1$	Λ_{n-1}	n
II	1A	3	0	$\{0\}$	0
II	1B	4	0	$\{0\}$	0
II	1C	6	0	$\{0\}$	0
II	1D	7	0	$\{0\}$	0
II	1E	9	0	$\{0\}$	0

continued on next page

continued from previous page

F	L	δ	r	Λ	d
II	1F	12	0	$\{0\}$	0
III	2	3	1	A_1	2
IV	3A	5	2	A_2	3
IV	3B	6	2	A_2	3
IV	3C	8	2	A_2	3
IV	3D	9	2	A_2	3
IV	3E	12	2	A_2	3
I_0^*	4A	6	4	D_4	4
I_0^*	4B	6	4	D_4	4
I_1^*	5A	7	5	D_5	4
I_2^*	5B	8	6	D_6	4
I_3^*	5C	9	7	D_7	4
I_4^*	5D	10	8	D_8	4
IV*	6A	9	6	E_6	3
IV*	6B	10	6	E_6	3
IV*	6C	12	6	E_6	3
III*	7	9	7	E_7	2
II*	8A	11	8	E_8	1
II*	8B	12	8	E_8	1

These invariants exclude certain configurations of singular fibres, as they must satisfy the conditions given in the following three lemmas.

Lemma A ([4]). *On a rational elliptic surface, $\sum \delta_F = 12$ and $\sum r_F \leq 8$.*

Lemma B ([5]). *If $\sum r_F = 8$, then $\prod d_F$ is a perfect square.*

Lemma C ([8]). *In order for a given configuration of singular fibres to exist, the associated lattice $\oplus_F \Lambda_F$ must embed in the root lattice E_8 .*

Oguiso and Shioda [6] provide a list, originally due to Dynkin [1], of lattices which embed in the root lattice E_8 . For the convenience of the reader we include the complement of that list here; namely, we list all direct sums of lattices of type A_n , D_n or E_n of rank no more than eight which do not embed in E_8 . Note that if the rank of such a lattice is 6 or less, it always embeds in E_8 .

Table 2: Lattices which are direct sums of A_n 's, D_n 's, or E_n 's, which do not embed in E_8 .

rank 7	rank 8
$A_2 \oplus A_1^{\oplus 5}$	$A_2 \oplus A_1^{\oplus 6}$ $A_2^{\oplus 2} \oplus A_1^{\oplus 4}$ $A_3 \oplus A_1^{\oplus 5}$

continued on next page

continued from previous page

rank 7	rank 8
$A_2^{\oplus 2} \oplus A_1^{\oplus 3}$	$A_3 \oplus A_2 \oplus A_1^{\oplus 3}$ $A_2^{\oplus 3} \oplus A_1^{\oplus 2}$
$A_3 \oplus A_2^{\oplus 2}$	$A_3 \oplus A_2^{\oplus 2} \oplus A_1$ $A_3^{\oplus 2} \oplus A_2$ $A_4 \oplus A_2^{\oplus 2}$ $A_5 \oplus A_3$ $A_6 \oplus A_2$
$A_4 \oplus A_1^{\oplus 3}$	$A_4 \oplus A_1^{\oplus 4}$ $A_4 \oplus A_2 \oplus A_1^{\oplus 2}$ $A_4 \oplus A_3 \oplus A_1$ $A_5 \oplus A_1^{\oplus 3}$ $A_6 \oplus A_1^{\oplus 2}$
$D_4 \oplus A_2 \oplus A_1$	$D_4 \oplus A_2 \oplus A_1^{\oplus 2}$ $D_4 \oplus A_2^{\oplus 2}$ $D_4 \oplus A_3 \oplus A_1$ $D_5 \oplus A_2 \oplus A_1$ $D_6 \oplus A_2$ $D_7 \oplus A_1$
	$D_5 \oplus A_1^{\oplus 3}$ $E_6 \oplus A_1^{\oplus 2}$

We also note that we have not used the J-map in our classifications at all, although it was used in the classifications done in other characteristics. We do not use it here because there were only a few cases where the J-map provided simple results, and each of these cases could be ruled out using another lemma which was required for more cases.

3 Surfaces with Singular Fibres of Multiplicative Type

We now consider the case where all singular fibres are of multiplicative type.

3.1 General Information

For convenience, when we write out a configuration of multiplicative fibres, we will write $n_1 n_2 n_3 \dots$ instead of writing $I_{n_1} I_{n_2} I_{n_3} \dots$. Repeated terms will be denoted by exponents, so $5 3^2 1$ denotes the configuration $I_5 I_3 I_3 I_1$.

As mentioned before, a rational elliptic surface has a multiplicative fibre over a point $t = p$ precisely if Δ has a root at p but b_2 does not. Since b_2 is

a non-zero polynomial of degree two, we have two cases: either b_2 has distinct roots, or b_2 has a repeated root. In all cases where a given configuration exists, we will give an example with distinct roots of b_2 . It is interesting to note that there are 11 cases which exist when b_2 has distinct roots but do not exist when b_2 has a repeated root; namely 531^4 , 81^4 , 52^21^3 , 541^3 , $3^22^21^2$, 62^21^2 , 5321^2 , 821^2 , 5^21^2 , 432^21 , and 4^22^2 .

We begin by reducing the discriminant using standard transformations to a normal form, given in equations (3.1.2) and (3.1.4). In the case that b_2 has distinct roots, we use a fractional linear transformation to move these roots to zero and infinity, and in the case that b_2 has a double root, we move that unique root to infinity.

Proposition 3.1.1. *Consider a minimal rational elliptic surface over $\mathbb{P}_t^1$ with Weierstrass equation*

$$y^2 = x^3 + b_2x^2 - b_4x + b_6, \quad (3.1.1)$$

where the b_i are polynomials in t of degree no more than i . If all of the singular fibres are of multiplicative type, and if the roots of b_2 are distinct, then up to automorphism of the surface and the base, the discriminant of the surface is a degree twelve polynomial of the form

$$\Delta = t^{12} + lt^{10} + t^3P_6(t) + nt^2 + m, \quad (3.1.2)$$

where $m^3 = n^2l^3 \neq 0$, and where $P_6(t)$ is a polynomial of degree 6.

Conversely, given any Δ of the form (3.1.2) there is a rational elliptic surface with Weierstrass equation (3.1.1) and with roots of b_2 at 0 and ∞ . If such a surface exists with Δ of form (3.1.2), we may, up to automorphism of the base, assume that one root of Δ occurs at $t = 1$.

Proof. $\text{PGL}(2)$ acts three-transitively on $\mathbb{P}^1$, thus we can find a transformation that sends one root of b_2 to infinity and one root to zero. Therefore, by a standard change of variables, we may assume $b_2 = t$.

Let

$$b_4 = \xi_4t^4 + \xi_3t^3 + \xi_2t^2 + \xi_1t + \xi_0,$$

and

$$b_6 = \gamma_6t^6 + \gamma_5t^5 + \gamma_4t^4 + \gamma_3t^3 + \gamma_2t^2 + \gamma_1t + \gamma_0,$$

where $\xi_i, \gamma_j \in k$. By substituting b_2, b_4, b_6 into the equation of Δ and simplifying we get

$$\begin{aligned} \Delta = & \xi_4^3t^{12} + \xi_4^2t^{10} + (-\gamma_6 - \xi_3\xi_4)t^9 + (\xi_3^2 - \gamma_5 - \xi_2\xi_4)t^8 \\ & + (-\gamma_4 - \xi_1\xi_4 - \xi_2\xi_3)t^7 + (-\gamma_3 + \xi_2^2 - \xi_0\xi_4 - \xi_1\xi_3 + \xi_2^2)t^6 \\ & + (-\gamma_2 - \xi_0\xi_3 - \xi_1\xi_2)t^5 + (-\gamma_1 - \xi_1^2 - \xi_0\xi_2^2)t^4 \\ & + (-\gamma_0 - \xi_0\xi_1 + \xi_1^3)t^3 + \xi_0^2t^2 + \xi_0^3. \end{aligned}$$

So

$$\Delta = a^3t^{12} + a^2t^{10} + t^3P_6 + b^2t^2 + b^3,$$

for some $a, b \in k$, and $P_6 \in k[t]$ is a polynomial of degree 6 in t . Since the surface has good reduction at infinity, we have $a \neq 0$; so using the transformation $x = a^{\frac{1}{2}}x'$ and $y = a^{\frac{3}{4}}y'$, we obtain

$$\Delta = (t^{12} + lt^{10} + t^3P_6 + mt^2 + n),$$

where

$$l = a^{-1}, \quad m = b^2a^{-3}, \quad n = b^3a^{-3},$$

and hence $m^3 = n^2l^3$.

Conversely, for any $l, m, n \in k$ such that $m^3 = n^2l^3 \neq 0$ and $P_6 \in k[t]$, we may invert the previous steps; first, by using the standard transformation $x' = a^{-\frac{1}{2}}x$ and $y' = a^{-\frac{3}{4}}y$ where $a = l^{-1}$, and $b = nm^{-1}$, giving $\Delta = a^3t^{12} + a^2t^{10} + t^3P_6 + b^2t^2 + b^3$. Letting

$$\xi_4 = a, \quad \xi_0 = b, \quad \epsilon_0 = \epsilon_2 = 0, \quad \epsilon_1 = 1$$

and $P_6(t) = \sum_{i=0}^6 \psi_i t^i$, with $\psi_i \in k$, we may choose any ξ_1, ξ_2 and $\xi_3 \in k$ and solve for the coefficients of b_6 :

$$\gamma_6 = -\xi_3\xi_4 - \psi_6, \quad \gamma_5 = \xi_3^2 - \xi_2\xi_4 - \psi_5, \quad \gamma_4 = -\xi_1\xi_4 - \xi_2\xi_3 - \psi_4,$$

$$\gamma_3 = \xi_2^3 - \xi_0\xi_4 - \xi_1\xi_3 + \xi_2^3 - \psi_3, \quad \gamma_2 = -\xi_0\xi_1 - \xi_1\xi_2 - \psi_2,$$

$$\gamma_1 = -\xi_1^2 - \xi_0\xi_2^3 - \psi_1, \quad \gamma_0 = -\xi_0\xi_1 + \xi_1^3 - \psi_0.$$

Substituting b_2, b_4 , and b_6 in Δ , we obtain

$$\Delta = a^3t^{12} + a^2t^{10} + t^3P_6 + b^2t^2 + b^3.$$

Finally, note that while $\text{PGL}(2)$ is 3-transitive, we only moved two points (the roots of b_2) to specific locations, so we may use $\text{PGL}(2)$ to move one root of Δ to $t = 1$ as well. $\square$

A similar argument proves the next proposition.

Proposition 3.1.2. *Consider a minimal rational elliptic surface over $\mathbb{P}_t^1$ with Weierstrass equation*

$$y^2 = x^3 + b_2x^2 - b_4x + b_6, \tag{3.1.3}$$

where b_i are polynomials in t and degree $b_i \leq i$. If all of the singular fibres are of multiplicative type, and if b_2 has a double root, then up to automorphism of the surface and the base, the discriminant Δ is a degree-twelve polynomial of the form

$$\Delta = t^{12} + lt^9 + mt^8 - nt^7 + P_6, \tag{3.1.4}$$

where $l, m, n \in k$, such that $n^3 = lm^3 \neq 0$, and $P_6(t)$ is a polynomial of degree less than or equal to 6. Conversely, given any Δ of the form 3.1.4 there is a rational elliptic surface with Weierstrass equation 3.1.3 and with roots of b_2 at ∞ . If such a surface exists, we may assume one root of Δ occurs at $t = 0$ and one occurs at $t = 1$.

To prove the existence of a rational elliptic surface with all singular fibres of multiplicative type, we simply need to exhibit a degree-twelve polynomial in the normal form with the zeros distributed in the desired way. Conversely, if no such polynomials exist, no surface with the prescribed fibre types exists. This yields the following lemma for non-existence of configurations with fibres of high multiplicity.

Lemma D. *A rational elliptic surface with purely multiplicative fibres cannot have three (not necessarily distinct) multiplicative fibres each of multiplicity at least three. In particular, no purely multiplicative configuration may include 3^3 , $3^2 4$, $3 4^2$, 4^3 , $3^2 5$, $3 4 5$, $3 6$, $3 7$, $3 8$, $4 6$, $5 6$, $5 7$, 6^2 , 9 , 10 , 11 , or 12 .*

Proof. Since we can move one root to $t = 1$, the discriminant Δ of such a surface can be written in the form:

$$\begin{aligned} (t-1)^3(t-c)^3(t-d)^3(t^3+et^2+ft+g) = & \quad (3.1.5) \\ t^{12} + et^{11} + ft^{10} + (g - c^3 - d^3 - 1)t^9 - (c^3e + d^3e + e)t^8 \\ & - (c^3f + d^3f - f)t^7 + (c^3 - d^3g - (1 + c^3)(g - d^3))t^6 \\ & + (c^3e + (1 + c^3)d^3e)t^5 + ((1 + c^3)d^3f + c^3f)t^4 \\ & + ((1 + c^3)d^3g + c^3(g - d))t^3 - c^3d^3et^2 - c^3d^3ft - c^3d^3g. \end{aligned}$$

If b_2 has distinct roots, Δ must be of the form

$$t^{12} + lt^{10} + t^3P_6^{(t)} + mt^2 + nb^3,$$

so the coefficient on the degree-11 term must vanish. Thus $e = 0$, and the polynomial is of the form

$$t^{12} + ft^{10} + t^3P_6 - c^3d^3ft - c^3d^3g.$$

Since the coefficient on the degree-two term must be nonzero to satisfy the form of Equation (3.1.2), we have a contradiction. Notice that no conditions were imposed on the distinctness of the roots of the polynomial.

Suppose now that b_2 has a double root. Since the coefficients of the degree-11 and degree-10 terms must be zero, we have

$$e = f = 0.$$

Thus we have

$$\Delta = t^{12} + (g - c^3 - d^3)t^9 + (c^3d^3 - d^3g - c^3g)t^6 + c^3d^3gt^3.$$

But since the coefficients of the degree-8 and degree-7 terms must be nonzero, no such polynomial satisfying Equation (3.1.4) exists. Again, no conditions were imposed on the distinctness of the roots. $\square$

3.2 Solutions for Purely Multiplicative Singularities

Below we list the 77 partitions of 12 using the notation of [4] and [2]. Next to each partition, we either list a polynomial of the form given in Equation (3.1.2) whose zeroes are distributed in the way indicated by the partition, or we indicate that the surface does not exist and give the letter of a lemma which proves its non-existence (often there are several that apply). As mentioned above, our results show that whenever a surface exists with purely multiplicative singularities and a double root of b_2 , one also exists with distinct roots of b_2 . We know of no good proof of this fact other than an appeal to our list below.

In the following list i denotes a root of $x^2 + 1 = 0$ in the field k .

1. **1¹²** $(t-1)(t^{11} + t^{10} - t^9 + t - 1) = t^{12} + t^{10} + t^9 + t^2 + t + 1$
2. **2¹1¹⁰** $(t-1)^2(t^{10} - t^9 + t^8 + t^2 + t - 1) = t^{12} + t^{10} + t^8 + t^4 - t^3 + t^2 - 1$
3. **3¹1⁹** $(t-1)^3(t^9 + t^7 + t^6 + t^3 - t^2 + 1) = t^{12} + t^{10} - t^7 - t^5 + t^2 - 1$
4. **2²1⁸** $(t^2 + 1)^2(t^8 - t^6 + t^4 - t^2 + 1) = t^{12} + t^{10} + t^2 + 1$
5. **4¹1⁸** $(t-1)^4(t^8 + t^7 - t^6 - t^3 - t^2 + t + 1) = t^{12} + t^{10} + t^7 - t^6 - t^5 + t^4 - t^3 + t^2 + 1$
6. **3²1⁷** $(t-1)^3(t+1)^2(t^7 + t^6 + t^5 - t^2 + t + 1) = t^{12} + t^{10} - t^9 + t^8 + t^7 - t^6 + t^5 + t^4 - t^3 + t^2 - 1$
7. **5¹1⁷** $(t-1)^5(t^7 - t^6 + t^5 + t^4 - t^2 + t - 1) = t^{12} + t^{10} - t^8 - t^7 - t^6 - t^3 + t^2 + 1$
8. **2³1⁶** $(t-1)^2(t^2 + 1)^2(t^6 - t^5 - t^4 + at^3 + t - 1)$
 $= t^{12} + t^{10} + (a+1)t^9 + (a+1)t^8 - at^6 + (a+1)t^4 + (a+1)t^3 + t^2 - 1$
where $a \neq 0, \pm 1$
9. **4²1⁶** $(t-1)^4(t+1)^2(t^6 - t^5 - t^2 + t - 1) = t^{12} + t^{10} - t^9 - t^7 + t^6 - t^5 + t^2 - 1$
10. **3²1⁶** $(t-1)^3(t+1)^3(t^6 + t^4 - t^2 + 1) = t^{12} + t^{10} - t^8 - t^4 + t^2 - 1$
11. **6¹1⁶** $(t-1)^6(t^6 + t^4 + t^2 + 1) = t^{12} + t^{10} + t^9 + t^8 + t^7 - t^6 + t^5 + t^4 + t^3 + t^2 + 1$
12. **3²2¹1⁵** $(t-1)^3(t^2 + 1)^2(t^5 - t^3 + 1) = t^{12} + t^{10} - t^9 - t^8 - t^6 - t^4 - t^3 + t^2 - 1$
13. **5²1⁵** $(t-1)^5(t+i+1)^2(t^5 + it^4 + t^3 + it^2 + (1+i)t + 1)$
 $= t^{12} + (i-1)t^{10} + it^9 + t^8 - (i+1)t^7 + (1+i)t^5 - t^4 + (i-1)t^3 + (1-i)t^2 + i$
14. **4³1⁵** $(t-1)^4(t+1)^3(t^5 + t^4 - t^3 - t - 1) = t^{12} + t^{10} + t^9 - t^8 - t^4 - t^3 + t^2 - 1$
15. **7¹1⁵** $(t-1)^7(t^5 + t^4 - t^3 + t^2 + dt + d) = t^{12} + t^{10} + (d-1)t^8 + t^7 - dt^6 + (d-1)t^5 + t^4 - (d+1)t^3 + (d-1)t^2 - d$
where $d^3 - d^2 - 1 = 0$
16. **2⁴1⁴** $(t-1)^2(t+1)^2(t^2 + 1)^2(t^4 + t^2 - 1) = t^{12} + t^{10} + t^6 + t^2 - 1$
17. **4²2¹1⁴** $(t-1)^4(t^2 + 1)^2(t^4 + t^3 + t^2 + t + 1) = t^{12} - t^{10} - t^9 + t^8 + t^4 - t^3 - t^2 + 1$

18. $\mathbf{3^2 2 1^4}$ $(t-1)^3(t+1)^3(t+i)^2(t^4+it^3+t^2+it-1) = t^{12}+t^{10}-it^9-t^8-t^4+it^3+t^2-1$
19. $\mathbf{6 2 1^4}$ $(t-1)^6(t+1)^2(t^4+t^3+t^2+t+1) = t^{12}+t^{10}-t^9+t^8+t^7+t^5+t^4-t^3+t^2+1$
20. $\mathbf{5 3 1^4}$ $(t-1)^5(t^3+1)(t^4-t^3+t^2-\alpha t+\alpha) = t^{12}+t^{10}-it^9+t^8+(i-1)t^6-t^4+it^3-t^2-i$
21. $\mathbf{4^2 1^4}$ $(t^4+1)^2(t^4+t^2-1) = t^{12}+t^{10}+t^8-t^6-t^4+t^2-1$
22. $\mathbf{8 1^4}$ $(t-1)^8(t^4-t^3+t^2+t-1) = t^{12}+t^{10}-t^9+t^8+t^7+t^6+t^5+t^4+t^2-1$
23. $\mathbf{3^2 3 1^3}$ $(t-1)^3(t+1)^2(t^2+it-1)^2(t^3+(1+i)t^2+(1-i)^2t+1-i) = t^{12}+t^{10}+t^9+t^8+(i-1)t^7+(1+i)t^6+(i-1)t^5-it^4+(1+i)t^3-it^2+i-1$
24. $\mathbf{5 2^2 1^3}$ $(t-1)^5(t^2+it+1)^2(t^3+(i-1)t^2+i(1-i)t-i) = t^{12}-it^{10}+t^9-(1+i)t^8-t^7-it^6+t^5+(i+1)t^4+t^3+it^2+i$
25. $\mathbf{4 3 2 1^3}$ $(t-1)^4(t+1)^3(t+i)^2(t^3+(1+i)t^2+(1-i)t+1) = t^{12}+it^{10}+t^9-it^8-it^4-t^3+it^2-1$
26. $\mathbf{7 2 1^3}$ $(t-1)^7(t+i)^2(t^3+(i+1)t^2+ct-c(i+1)) = t^{12}+(c-i-1)t^{10}+(ic+c)t^9+(1-c+i)t^8+(c-i-1)t^7+(1-c+i)t^5+(c-i-1)t^4-t^3+(1-c+i)t^2-c-ic$
where $ic^5-c^3-(i+1)c^2-i+1=0$
27. $\mathbf{3^3 1^3}$ does not exist, D
28. $\mathbf{6 3 1^3}$ does not exist, D
29. $\mathbf{5 4 1^3}$ $((t-1)^5(t+i)^4(t^3-(1+i)t^2+t-i-1)) = t^{12}-(i-1)t^{10}+t^9-(i+1)t^8-it^7+(1-i)t^6-it^5+(1-i)t^4+(1-i)t^3+(1-i)t^2-1-i$
30. $\mathbf{9 1^3}$ does not exist, D
31. $\mathbf{2^5 1^2}$ $(t^5-1)^2(t^2+1) = t^{12}+t^{10}+t^7+t^5+t^2+1$
32. $\mathbf{4 2^3 1^2}$ $(t-1)^4(t^2+t+1)^2(t+1)^2(t^2+1) = t^{12}-t^{10}+t^9-t^8-t^7-t^6-t^5-t^4+t^3-t^2+1$
33. $\mathbf{3^2 2^2 1^2}$ $(t-1)^3(t+1)^3(t^2+i)^2(t^2+i+1) = t^{12}+t^{10}-it^8+(1-i)t^6-t^4+it^2+i+1$
34. $\mathbf{6 2^2 1^2}$ $(t-1)^6(t^2+1)^2(t^2+i) = t^{12}+(i-1)t^{10}+t^9+(1-i)t^8+(i-1)t^7+(i+1)t^6+(1-i)t^5+(i-1)t^4+it^3+(1-i)t^2+i$
35. $\mathbf{5 3 2 1^2}$ $(t-1)^5(t^3+i)(t+1)^2(t^2-1-i) = t^{12}-it^{10}+(i-1)t^9-it^8+(1+i)t^7+(i-1)t^6+(1+i)t^5-t^4-t^3-t^2+i-1$
36. $\mathbf{4^2 2 1^2}$ $(t-1)^4(t-i)^4(t+1-i)^2(t^2-t+i) = t^{12}+t^{10}+it^9+it^8+(i-1)t^7-(1+i)t^6-(1+i)t^5-it^4+t^3+t^2-1$

37. **8 2 1²** $(t-1)^8(t+1)^2(t^2+1) = t^{12}-t^{10}+t^9-t^8-t^7-t^6-t^5-t^4+t^3-t^2+1$
38. **4 3² 1²** does not exist, C
39. **7 3 1²** does not exist, B
40. **6 4 1²** does not exist, C
41. **5² 1²** $(t-1)^5(t+1)^5(t^2+i)$
 $= t^{12} + (1+i)t^{10} + (1+i)t^8 + (i-1)t^6 - (1+i)t^4 - (1+i)t^2 - i$
42. **10 1²** does not exist, A
43. **3 2⁴ 1** $(t-1)^3(t^4+t^3-t^2-t-1)^2(t+1) = t^{12}+t^{10}-t^8-t^7-t^6+t^3+t^2-1$
44. **5 2³ 1** does not exist, C
45. **4 3 2² 1** $(t-1)^4(t^3-i)(t^2+c)^2(t+1)$
 $= t^{12} - (1+c)t^{10} + (i-1)t^9 + (c^2+c)t^8 + (c-i+1-ic)t^7 - (c^2+i)t^6 +$
 $(ic^2-c^2+ic-c)t^5 + (ic+i)t^4 + (c^2-ic^2)t^3 - (ic+ic^2)t^2 + ic^2$
where $c^4 - ic^3 + c - i = 0$
46. **7 2² 1** does not exist, B
47. **3³ 2 1** does not exist, D
48. **6 3 2 1** does not exist, D
49. **5 4 2 1** does not exist, B
50. **9 2 1** does not exist, A
51. **5 3² 1** does not exist, C
52. **4² 3 1** does not exist, C
53. **8 3 1** does not exist, A
54. **7 4 1** does not exist, A
55. **6 5 1** does not exist, A
56. **11 1** does not exist, A
57. **2⁶** $(t-1)^2(t^5+t^4+t+1)^2 = t^{12}+t^{10}-t^6+t^2+1$
58. **4 2⁴** $(t-1)^4(t^4-t^3+t^2-t+1)^2 = t^{12}-t^{10}+t^9+t^7-t^6+t^5+t^3-t^2+1$
59. **3² 2³** does not exist, C
60. **6 2³** does not exist, C
61. **5 3 2²** does not exist, C
62. **4² 2²** $(t-1)^4(t+1)^4(t^2+1)^2 = t^{12}+t^{10}-t^8+t^6-t^4+t^2+1$
63. **8 2²** does not exist, A
64. **4 3² 2** does not exist, B

- 65. **7 3 2** does not exist, A
- 66. **6 4 2** does not exist, A
- 67. **5² 2** does not exist, A
- 68. **10 2** does not exist, A
- 69. **3⁴** does not exist, D
- 70. **6 3²** does not exist, A
- 71. **5 4 3** does not exist, A
- 72. **9 3** does not exist, A
- 73. **4³** does not exist, A
- 74. **8 4** does not exist, A
- 75. **7 5** does not exist, A
- 76. **6²** does not exist, A
- 77. **12** does not exist, A

4 At Least One Additive Fibre

We now consider the case where the rational elliptic surface has at least one additive fibre.

4.1 General Information

In each of the cases below we will use a linear transformation to send the worst singularity to $t = 0$, using the fact that $\mathrm{PGL}(2)$ acts three-transitively on $\mathbb{P}^1$. We classify “worst” singularities using the following criteria:

1. Additive type singularities are worse than multiplicative type singularities.
2. If comparing two additive type singularities with different values of δ (where δ is the multiplicity of the root of Δ over which the singularity happens), the singularity with the largest δ is worse.
3. If two additive type singularities have the same value of δ , the singularity with the largest r value is worse.

As in [3], we may change coordinates so that the Weierstrass equation becomes

$$y^2 = x^3 + tc_1x^2 + tc_3x + tc_5, \quad (4.1.1)$$

where the c_i are polynomials in t of degree no more than i . In our notation,

$$b_2 = tc_1, \quad b_4 = -tc_3, \quad \text{and} \quad b_6 = tc_5,$$

and Δ becomes

$$-\Delta = t^4 c_1^2 (c_1 c_5 - c_3^2) + t^3 c_3^3. \quad (4.1.2)$$

A list of all singular fibres of additive type that could potentially appear on a rational elliptic surface in characteristic three was given in [3], and a normal form over $t = 0$ was given for each one. We will use the normal forms of [3] for each of the worst singularities. For the reader's convenience, these normal forms are listed at the beginning of each case treated in Section 4.2.

An easy application of Tate's algorithm (or a glance at Equation (4.1.2)) shows that all additive fibres have $\delta \geq 3$. The following lemma is also an easy result of Tate's algorithm [9].

Lemma E. *If $b_2 \equiv 0$, then all singular fibres on the surface are of additive type. If $t^2|b_2$, where b_2 is not identically zero, then there is only one additive singular fibre.*

Since $\text{PGL}(2)$ is three-transitive, after sending the worst singularity to $t = 0$, we may still move two other points of the base—either singular fibres or roots of b_2 .

As in the purely multiplicative situation, we may consider two cases: when b_2 has a double root at $t = 0$, and when b_2 has distinct roots. If b_2 has distinct roots, we can move the second root of b_2 to $t = \infty$. Again, this gives some significant restrictions on Δ .

Proposition 4.1.1. *Consider a minimal rational elliptic surface over $\mathbb{P}_t^1$ with a singular fibre of additive type over $t = 0$ and Weierstrass equation*

$$y^2 = x^3 + tc_1x^2 + tc_3x + tc_5, \quad (4.1.3)$$

where the c_i are polynomials in t of degree no more than i . If the roots of b_2 are distinct, then up to automorphism of the surface and the base, the discriminant of the surface is a polynomial of degree 12 or less, of the form

$$\Delta = -\alpha^3 t^{12} + \alpha^2 \beta^2 t^{10} + t^6 P_3(t) + \beta^2 t^4 P_1(t) - \sigma^3 t^3, \quad (4.1.4)$$

where α , β , and σ are elements of k , and the $P_i(t)$ are elements of $k[t]$ of degree no more than i . In particular, the t^{11} term vanishes.

Similarly, if b_2 has a double root, then up to automorphism of the surface and the base, the discriminant of the surface is a polynomial of degree 12 or less, of the form

$$\Delta = t^6 P_6(t) - \sigma t^3, \quad (4.1.5)$$

where σ is an element of k , and P_6 is an element of $k[t]$ of degree no more than 6. In particular, the t^4 and t^5 terms vanish.

Finally, if b_2 is identically zero, then up to automorphism of the surface and the base, the discriminant of the surface is a polynomial of degree 12 or less, of the form

$$\Delta = b_4^3 = -t^3 (c_3)^3. \quad (4.1.6)$$

Proof. We let

$$\begin{aligned} b_2 = tc_1 &= t(\epsilon_1 t + \epsilon_0) \\ b_4 = -tc_3 &= t(\phi_3 t^3 + \phi_2 t^2 + \phi_3 t + \phi_0) \\ b_6 = tc_5 &= t(\gamma_5 t^5 + \gamma_4 t^4 + \cdots + \gamma_0), \end{aligned} \quad (4.1.7)$$

with ϵ_i , ϕ_i and γ_i in k . If b_2 has distinct roots, then we may move the second root to infinity, which amounts to setting $\epsilon_1 = 0$. We substitute these into the expression (4.1.2) for Δ to get

$$\begin{aligned} \Delta = & -\phi_3^3 t^{12} + \epsilon_0^2 \phi_3^2 t^{10} - (\epsilon_0^2 \phi_3 \phi_2 + \epsilon_0^3 \gamma_5 + \phi_2^3) t^9 \\ & + (2\epsilon_0^3 \gamma_4 + \epsilon_0^2 \phi_2^2 + 2\epsilon_0^2 \phi_3 \phi_1) t^8 \\ & - (\epsilon_0^3 \gamma_3 + \epsilon_0^2 \phi_2 \phi_1 + \epsilon_0^2 \phi_3 \phi_0) t^7 \\ & + (\epsilon_0^2 \phi_1^2 + 2\phi_1^3 + 2\epsilon_0^2 \phi_2 \phi_0 + 2\epsilon_0^3 \gamma_2) t^6 \\ & + (2\epsilon_0^3 \gamma_1 + 2\epsilon_0^2 \phi_1 \phi_0) t^5 + (2\epsilon_0^3 \gamma_0 + \epsilon_0^2 \phi_0^2) t^4 + 2t^3 \phi_0^3. \end{aligned} \quad (4.1.8)$$

This has the desired form.

Similarly, if b_2 has a double root, we may assume that $\epsilon_0 = 0$, and substituting into Equation (4.1.2) gives

$$\begin{aligned} \Delta = & (\epsilon_1^2 \phi_3^2 + 2\epsilon_1^3 \gamma_5 + 2\phi_3^3) t^{12} + (2\epsilon_1^3 \gamma_4 + 2\epsilon_1^2 \phi_3 \phi_2) t^{11} \\ & + (2\epsilon_1^2 \phi_3 \phi_1 + \epsilon_1^2 \phi_2^2 + 2\epsilon_1^3 \gamma_3) t^{10} \\ & + (2\epsilon_1^2 \phi_2 \phi_1 + 2\epsilon_1^2 \phi_3 \phi_0 + 2\epsilon_1^3 \gamma_2 + 2\phi_2^3) t^9 \\ & + (2\epsilon_1^2 \phi_2 \phi_0 + 2\epsilon_1^3 \gamma_1 + \epsilon_1^2 \phi_1^2) t^8 \\ & + (2\epsilon_1^3 \gamma_0 + 2\epsilon_1^2 \phi_1 \phi_0) t^7 + (\epsilon_1^2 \phi_0^2 + 2\phi_1^3) t^6 + 2t^3 \phi_0^3. \end{aligned} \quad (4.1.9)$$

This again has the desired form.

Finally, substituting $b_2 = 0$ into Equation (4.1.2) gives the last form. $\square$

An immediate corollary of the proposition is the following lemma.

Lemma F. *The configuration $\mathbf{X}3^221$ is not possible, where $\mathbf{X}$ is any additive fibre. Nor is any specialization of this configuration possible, provided the $\mathbf{I}_2$ and $\mathbf{I}_1$ fibres remain distinct from each other and both the order-three multiplicative singularities remain distinct from the additive fibre $\mathbf{X}$. More precisely, the following configurations cannot exist: $\mathbf{X}3^221$, $\mathbf{X}621$, $\mathbf{X}531$, $\mathbf{X}432$, $\mathbf{X}54$, $\mathbf{X}3^21$, $\mathbf{X}61$, $\mathbf{X}3^22$, $\mathbf{X}62$, $\mathbf{X}43$, $\mathbf{X}53$.*

Proof. Assume this configuration exists. $\mathbf{X}$ must have $\delta = 3$, since $\sum \delta_i = 12$, and we may assume $\mathbf{X}$ occurs over $t = 0$. We may further assume no singular fibre occurs over infinity. We let the first $\mathbf{I}_3$ occur over $t = a$, the second $\mathbf{I}_3$ over $t = b$, the $\mathbf{I}_2$ over $t = c$, and the $\mathbf{I}_1$ over $t = d$.

The hypotheses of the lemma require $c \neq d$, and $ab \neq 0$ but do not otherwise require that the roots be distinct or non-vanishing. We have $\eta t^3(t - a)^3(t -$

$b)^3(t-c)^2(t-d) = \Delta$ for some $\eta \in k^*$. We expand to get

$$\begin{aligned} t^3(t-a)^3(t-b)^3(t-c)^2(t-d) = & \\ & t^{12} + (c+2d)t^{11} + (2cd+c^2)t^{10} \\ & + (2b^3+2c^2d+2a^3)t^9 + (2a^3c+2b^3c+a^3d+b^3d)t^8 \\ & + (2b^3c^2+a^3cd+b^3cd+2a^3c^2)t^7 \\ & + (a^3b^3+a^3c^2d+b^3c^2d)t^6 + (2a^3b^3d+a^3b^3c)t^5 \\ & + (2a^3b^3cd+a^3b^3c^2)t^4 + 2t^3a^3b^3c^2d. \end{aligned}$$

If b_2 has distinct roots, the t^{11} term of Δ must vanish, as in equation (4.1.4), which means $c = d$, a contradiction.

If b_2 has a repeated root, then the t^5 term of Δ must vanish, as in equation (4.1.5), which means $a^3b^3c = a^3b^3d$, and thus either $c = d$ or $ab = 0$, a contradiction. $\square$

All of the configurations that do not exist fail to do so by one of the previous lemmas, or by straightforward algebraic manipulations. We list all that fail from algebraic manipulations in the following lemma.

Lemma G. *The following configurations do not exist:*

1. **II 3²** and **II 6**.
2. **IV 2³** and **IV 4 2**.
3. **IV 3 2 1** and **IV 5 1**.
4. **I₁^{*} 4 1**.
5. **III 4² 1** and **III 5 4**.
6. **IIII 3²**, **IIII 6**, **IIII 3²**, and **IIII 6**.
7. **IIII 4 2**, and **IIII 4 2**, where the I_4 and I_2 are distinct.

Proof. 1. **II 3²** and **II 6**. Suppose this configuration exists. Assume that one I_3 singularity occurs over $t = a$ and the other over $t = b$. We allow the possibility of $a = b$. The fibre of type II must have $\delta = 6$, which means it is case 1C, which requires that $\epsilon_0 = 0$, $\phi_0 = 0$, and $\gamma_0 \neq 0$ (see Section 4.2.3). Making the substitutions of Equation (4.1.7) for b_i in the discriminant, we have

$$\begin{aligned} \alpha[(t-a)^3(t-b)^3] &= -\Delta/t^6 \\ &= (2\phi_3^3 + 2\epsilon_1^3\gamma_5 + \epsilon_1^2\phi_3^2)t^6 + (2\epsilon_1^2\phi_3\phi_2 + 2\epsilon_1^3\gamma_4)t^5 \\ &\quad + (\epsilon_1^2\phi_2^2 + 2\epsilon_1^3\gamma_3 + 2\epsilon_1^2\phi_3\phi_1)t^4 \\ &\quad + (2\epsilon_1^3\gamma_2 + 2\epsilon_1^2\phi_2\phi_1 + 2\phi_2^3)t^3 \\ &\quad + (\epsilon_1^2\phi_1^2 + 2\epsilon_1^3\gamma_1)t^2 + 2t\epsilon_1^3\gamma_0 + 2\phi_1^3, \end{aligned}$$

for some $\alpha \in k$.

Since the right-hand side of the above equation has a t term, but the left-hand side does not, the t term must vanish. But $\epsilon_1 = 0$ would force $b_2 = 0$, and thus all fibres would be additive, and $\gamma_0 = 0$ is prohibited in case 1C.

2. **IV 2³** and **IV 4 2**. Suppose this configuration exists. Since the type **IV** fibre occurs with $\delta = 6$, we are in case 3C, with $\epsilon_1 = 0$, $\phi_0 = 0$, $\gamma_0 = 0$, but $\phi_1 \neq 0$. Assume the multiplicative fibres occur over a , b , and c in $\mathbb{A}^1 - \{0\}$. If all $a = b = c$, we have the configuration **IV 6**, which is possible (see Section 4.2.9). If any two of these fibres are distinct, we may assume, without loss of generality, that $a = 1$ and $b = -1$, which means we have

$$\begin{aligned} \alpha[t^6 + ct^5 + (1 + c^2)t^4 + ct^3 + (1 + c^2)t^2 + ct + c^2] &= -\Delta/t^6 \\ &= (-\epsilon_1^3\gamma_5 + \epsilon_1^2\phi_3^2 - \phi_3^3)t^6 + (2\epsilon_1^3\gamma_4 - \epsilon_1^2\phi_3\phi_2)t^5 + (\epsilon_1^2\phi_2^2 - \epsilon_1^2\phi_3\phi_1 - \epsilon_1^3\gamma_3)t^4 \\ &\quad + (2\phi_2^3 - \epsilon_1^2\phi_2\phi_1 - \epsilon_1^3\gamma_2)t^3 + (2\epsilon_1^3\gamma_1 + \epsilon_1^2\phi_1^2)t^2 - \phi_1^3. \end{aligned}$$

Comparing low-degree coefficients shows that $c = 0$, a contradiction.

3. **IV 3 2 1**, **IV 5 1**. Suppose this configuration exists. Again the **IV** falls in case 3C, and we assume the I_3 fibre occurs over a , the I_2 over b , and the I_1 over c , with $abc \neq 0$. If the I_2 and I_1 fibres are distinct, we have $b \neq c$, and we may assume that $b = 1$ and $c = -1$, which yields the following:

$$\begin{aligned} \alpha[t^6 + 2t^5 + 2t^4 + (1 + 2a^3)t^3 + a^3t^2 + a^3t + 2a^3] &= -\Delta/t^6 \\ &= (-\epsilon_1^3\gamma_5 + \epsilon_1^2\phi_3^2 - \phi_3^3)t^6 + (2\epsilon_1^3\gamma_4 - \epsilon_1^2\phi_3\phi_2)t^5 + (\epsilon_1^2\phi_2^2 - \epsilon_1^2\phi_3\phi_1 - \epsilon_1^3\gamma_3)t^4 \\ &\quad + (2\phi_2^3 - \epsilon_1^2\phi_2\phi_1 - \epsilon_1^3\gamma_2)t^3 + (2\epsilon_1^3\gamma_1 + \epsilon_1^2\phi_1^2)t^2 - \phi_1^3. \end{aligned}$$

Comparing low-degree coefficients shows that $a = 0$, a contradiction.

4. **I₁^{*} 4 1**. Suppose this configuration exists. The **I₁^{*}** fibre falls into case 5A, and thus has $\phi_0 = \phi_1 = \phi_2 = 0$ and $\gamma_0 = \gamma_1 = \gamma_2 = 0$, but $\epsilon_0 \neq 0$ and $\gamma_3 \neq 0$. In particular, b_2 has distinct roots. We may assume that the non-zero root of b_2 occurs at infinity, which implies that $\epsilon_1 = 0$, and the I_4 fibre occurs at $t = 1$. The remaining I_1 fibre occurs over some $a \in \mathbb{A}^1 - \{0, 1, \infty\}$. Making these substitutions, we have

$$\begin{aligned} \alpha[t^5 - (1 + a)t^4 + t^3a - t^2 + (1 + a)t - a] &= \Delta/t^7 \\ &= 2\phi_3^3t^5 + \phi_3^2t^3 - t^2\gamma_5 + 2t\gamma_4 + 2\gamma_3. \end{aligned}$$

Since the coefficient of t^4 vanishes in Δ/t^7 , we have that $a = 1$, which is a contradiction.

5. **III 4² 1** and **III 5 4**. Suppose this configuration exists. The type **III** fibre falls into case 2, which means that $\gamma_0 = 0$, but $\phi_0 \neq 0$ and $\epsilon_0 \neq 0$. Let

the I_4 's happen at $t = a$ and $t = b$, and let the I_1 happen at $t = c$. We require that $a \neq b$, but we could have $a = c$ or $b = c$. We have

$$\Delta = \alpha(t^3(t-a)^4(t-b)^4(t-c)) \quad (4.1.10)$$

for some $\alpha \neq 0$. If b_2 has two distinct roots, then by Proposition 4.1.1 we may assume that $\epsilon_1 = 0$ and Δ has no t^{11} term. This means $c = -a - b$. Furthermore, we may assume that $a = 1$. Making these substitutions into Δ shows that the leading term vanishes if ϕ_3 does, which would put an additional additive singularity at infinity. Thus ϕ_3 is non-zero. Equating like terms in Equation (4.1.10), we obtain the following:

$$\begin{aligned} \frac{\phi_0^3}{\phi_3^3} &= b^5 + b^4 & (\text{from } t^3) \\ -\frac{\epsilon_0^2 \phi_0^2}{\phi_3^3} &= -b^4 - b^3 - b^5 & (\text{from } t^4) \\ -\frac{\epsilon_0^2}{\phi_3} &= -1 - b - b^2 & (\text{from } t^{10}). \end{aligned}$$

A straightforward manipulation of these constraints gives $b = 1$, which is a contradiction, since $a = 1$ and a and b are distinct.

If b_2 has a repeated root, by Proposition 4.1.1 the t^5 and t^4 terms must vanish, and we may assume that $a = 1$ and $b = -1$. Substituting into Equation (4.1.10), and setting like terms equal, we get that $c = 0$, which is a contradiction.

6. **IIIIII3², IIIIII6, IIIII3², and IIIII6.**

Suppose this configuration is possible. We may assume that the type **III** fibre occurs over $t = 0$, which implies that $\phi_0 \neq 0$, but $\gamma_0 = 0$. We further assume that the other additive fibre occurs over $t = \infty$, so that b_2 has distinct roots, and we may take $\epsilon_0 = 1$ and $\epsilon_1 = 0$. Furthermore, suppose the first I_3 happens at $t = a$ for some $a \neq 0, \infty$, and that the second I_3 happens at $t = b \neq 0, \infty$. Making these substitutions into Δ shows that the coefficient of t^4 in Δ is ϕ_0^2 . But matching coefficients in the equality

$$\Delta = \alpha t^3(t-a)^3(t-b)^3 = t^9 - (a^3 + b^3)t^6 - t^3 a^3 b^3,$$

with $\alpha \in k^*$, shows that $\phi_0 = 0$, which is a contradiction.

7. **IIIIII4 2 and IIIII4 2.** Suppose this configuration is possible. As in the previous case, we may assume that $\gamma_0 = \epsilon_1 = 0$ and $\epsilon_0 = 1$. Since the t^8 term vanishes in the expansion of Δ , matching coefficients in the equality

$$\Delta = \alpha(t-a)^4(t-b)^2$$

shows that $a = b$, which is a contradiction. □

4.2 Solutions for Additive and Multiplicative Singularities

We list Lang's type [3] for the worst singularity, its main invariants, Weierstrass form, and $-\Delta$. For each subsequent configuration, we list the Weierstrass form and $-\Delta$ if the desired rational elliptic surface exists, or the lemma number proving impossibility if it does not exist. In the following examples, i denotes a root of $x^2 + 1 = 0$ in k .

4.2.1 Case 1A

$$\delta = 3, \quad r = 0 \quad \text{Type II}$$

$$\begin{aligned} y^2 &= x^3 + tc_1x^2 + tc_3x + tc_5, \quad t \nmid c_5, t \nmid c_3 \\ -\Delta &= t^2c_1^2(t^2c_1c_5 - t^2c_3^2) + t^3c_3^3 \end{aligned} \quad (4.2.1)$$

1. **II 9** $y^2 = x^3 + t^2x^2 + t(t+1)x + t(t+2)$
 $-\Delta = 2t^3$
2. **II 8 1** does not exist, F
3. **II 7 2** does not exist, F
4. **II 7 1²** $y^2 = x^3 + t(t+i)x^2 + t(2t+i)x + t^2$
 $-\Delta = -2it^3 + t^4 - 2it^5 = -t^3(2+it+2t^2)i$
5. **II 6 3** $y^2 = x^3 + t^2x^2 + t(2t+2)x + t(2+t)$
 $-\Delta = t^9 + 2t^6 = t^6(t^3+2)$
6. **II 6 2 1** does not exist, F
7. **II 6 1³** $y^2 = x^3 + t(t+1)x^2 + t(t+2+i)x + t(t+2i)$
 $-\Delta = (-1-i)t^6 + it^5 - it^4 + (1+i)t^3 = -t^3(t-1)((1+i)t^2 + t + (1+i))$
8. **II 5 4** does not exist, F
9. **II 5 3 1** does not exist, F
10. **II 5 2²** $y^2 = x^3 + t\epsilon_0x^2 + t(2t^3+2+i)x + \frac{t}{\epsilon_0}((2+2i)t^5 + 2it^4 + (2+i)t^3 + t^2 + (2+2i)t + 2i)$
 $-\Delta = t^{12} + it^{10} + (-1+i)t^9 + 2t^8 + 2it^6 + (-1+i)t^5 + t^4 + (1+i)t^3 = t^3(t+(1+i))^5(t-1)^2(t-i)^2$, where $\epsilon_0^2 = i$
11. **II 5 2 1²** $y^2 = x^3 + tx^2 + t(2t^3+i)x + t(it^5 + t^4 + 2it^3 + t^2 + 2it + 2)$
 $-\Delta = t^{12} + t^{10} + 2it^9 + 2t^8 + 2it^7 + 2t^6 + it^5 + it^3 = t^3(t+i)^5(t-i)^2(t-1)(t+1)$
12. **II 5 1⁴** $y^2 = x^3 + t\epsilon_0x^2 + t(2t^3+i)x + \frac{t}{\epsilon_0}((1-i)t^5 + (1+i)t^4 + t^3 + (1+i)t^2 + it - i)$
 $-\Delta = t^{12} + (1+i)t^{10} + t^9 + it^8 + t^7 + it^6 + (1+2i)t^5 + t^4 + it^3 = t^3(t-(1-i))^5(t-i)(t+1)(t-1)(t-(1+i))$

13. **II 4² 1** $y^2 = x^3 + t(t + \epsilon)x^2 + tx + t\left(\frac{-t}{\epsilon-1} + \frac{\epsilon}{\epsilon-1}\right)$
 $-\Delta = \frac{t^8}{\epsilon-1} - \frac{\epsilon t^7}{\epsilon-1} + t^6 - \frac{t^5}{\epsilon-1} + \frac{\epsilon t^4}{\epsilon-1} + \frac{(1-\epsilon)t^3}{\epsilon-1} = t^3(t-1)^4\left(2 + \frac{1}{\epsilon-1} - \frac{\epsilon}{\epsilon-1} - 1 + \epsilon^2\right)t$,
where $\epsilon^3 = \epsilon^2 - \epsilon - 1$
14. **II 4 3 2** does not exist, F
15. **II 4 3 1²** $y^2 = x^3 + t(t+1)x^2 + tx + t(t+2)$
 $-\Delta = 2t^8 + t^7 + t^6 + t^5 + 2t^4 + 2t^3 = -t^3(t-1)^3(2-t+t^2)$
16. **II 4 2² 1** $y^2 = x^3 + tx^2 + t(2t^3 + 2i)x + t(2t^5 + it^4 - it^2 + t + i)$
 $-\Delta = t^{12} + 2t^{10} - it^9 + 2t^8 + it^7 + t^6 + it^5 + 2it^3 = t^3(t+i)^4(t-1)^2(t+1)^2(t-i)$
17. **II 4 2 1³** $y^2 = x^3 + t\epsilon x^2 + t(2t^3 + 1)x + \frac{t}{\epsilon}(-t^5 - (1+i)t^4 - t^3 + (1-i)t^2 + t - 1)$
 $-\Delta = t^{12} + (2-i)t^{10} + (2+2i)t^9 + it^8 + (1+i)t^7 + 2t^6 + (1+i)t^5 + (1+i)t^4 + 2t^3 = t^3(t + (1+i))^4(t+i)^2(t - (1+i))(t - (1-i))(t+1)$, where $\epsilon^2 = 2 - i$
18. **II 4 1⁵** $y^2 = x^3 + t\epsilon x^2 + t(2t^3 + 1 + i)x + \frac{t}{\epsilon}((1-2i)t^5 - it^4 + t^2 + (2-i)t - 2i)$
 $-\Delta = t^{12} + it^{10} + (1-i)t^9 + 2t^8 + (2+i)t^7 - it^6 + (2+i)t^5 + 2t^4 + (2+i)t^3 = t^3(t - (2-i))^4(t-1)(t+i)(t-i)(t+1)(t - (1+i))$, where $\epsilon^2 = i$
19. **II 3³** $y^2 = x^3 + t^2x^2 + t(t+1)x + t(2+2t^2+t)$
 $-\Delta = t^9 + 2t^3 = t^3(t-1)^3(t+1)^3$
20. **II 3² 2 1** does not exist, F
21. **II 3² 1³** $y^2 = x^3 + t(t+1)x^2 + t(2t+1)x + t(2+2t^2+2t)$
 $-\Delta = t^9 + 2t^8 + t^7 + t^5 + 2t^4 + 2t^3 = -t^3(t-1)^3(-t^3 + t^2 - t - 1)$
22. **II 3 2³** $y^2 = x^3 + t\epsilon x^2 + t(2t^3 + i)x + \frac{t}{\epsilon}((1-i)t^5 - it^4 - 2it^2 - it - i)$
 $-\Delta = t^{12} + it^{10} + (2+2i)t^9 + 2t^8 + 2t^7 + t^6 + 2t^5 + (2-i)t^4 - 2it^3 = t^3(t+1)^3(t - (2-i))^2(t-1)^2(t-i)^2$, where $\epsilon^2 = i$
23. **II 3 2² 1²** $y^2 = x^3 + t(t+1)x^2 + t(t+2)x + t((2+i)t^2 + (2+i)t + 2+i)$
 $-\Delta = (1-i)t^9 - (1+i)t^8 + (1-i)t^7 + (1-i)t^6 + (1-i)t^5 - (1+i)t^4 + t^3 = i\left(t - \frac{1}{1-i}\right)^2(2t^2 + 2it^2 + 1)t^3(t-1)^2$
24. **II 3 2 1⁴** $y^2 = x^3 + t^2x^2 + t(t+1)x + t(1+t^2)$
 $-\Delta = 2t^9 + t^8 + t^7 + 2t^3 = t^3(t-1)^2(-t^4 - t^3 + t - 1)$
25. **II 3 1⁶** $y^2 = x^3 + (t^2 + it)x^2 + (it^2 + 2t + t^2)x + 2t^3 + 2t^2 - it^2 + it$
 $-\Delta = t^9 + t^8 + t^6 + t^4 + t^3 = t^3(t^6 + t^5 + t^3 + t + 1)$
26. **II 2⁴ 1** $y^2 = x^3 + t^2x^2 + 2tx + t(2t^5 + 2t^4 + 1 + t^2 + 2t)$
 $-\Delta = t^{12} + t^{11} + 2t^9 + t^8 + 2t^7 + t^6 + t^3 = t^3(t-1)^2(t^3 + t^2 + t + 2)^2(t-2)$
27. **II 2³ 1³** $y^2 = x^3 + itx^2 + t(2t^3 + 2 + i)x + t((1+2i)t^5 + 2 + (2+i)t^3 + 2t^2 + 2it)$
 $-\Delta = t^{12} + 2t^{10} + (1+i)t^9 + it^7 - it^6 + t^5 + it^4 + (1+i)t^3 = t^3(t - (1+i))^2(t + (1+i))^2(t+1)^2(t-1)(t-i)(t - (1-i))$

28. **II 2² 1⁵** $y^2 = x^3 + t\epsilon x^2 + t(2t^3 + 2i)x + \frac{t}{\epsilon}((1-i)t^5 - it^4 + t^3 + t^2 + (-1+i)t + (1-i))$
 $-\Delta = t^{12} + (1+i)t^{10} + t^9 + (2+i)t^8 + it^7 + (2+2i)t^6 + 2t^5 + 2it^4 + 2it^3t^3(t-(1+i))^2(t-1)^2(t+1)(t+i)(t-i)(t+(1+i))(t-(1-i))$, where $\epsilon^2 = 1+i$
29. **II 2¹ 1⁷** $y^2 = x^3 + t\epsilon x^2 + t(2t^3 + 2+i)x + \frac{t}{\epsilon}((-1+2i)t^5 + it^4 + (-2+2i)t^3 + t^2 + (-2+i)t + i)$
 $-\Delta = t^{12} - it^{10} + (4+2i)t^9 + 2t^8 + (5+5i)t^7 - 2it^6 + (2+4i)t^5 + (4+4i)t^3 = t^3(t-(-1-i))^2(t-1)(t+1)(t-i)(t-(-1-i))(t+i)(t-(-1+i))(t-(1-i))$, where $\epsilon^2 = -i$
30. **II 1⁹** $y^2 = x^3 + t(t+1)x^2 + 2tx + t(t^4 + 2t^3 + 1)$
 $-\Delta = 2t^{11} + t^{10} + 2t^8 + t^6 + 2t^5 + t^3 = -t^3(t-1)(1+t+t^3+t^4+t^7)$
31. **II III III** $y^2 = x^3 - t(t^2 - 1)x + t(t^2 - 1)(t^2 - t - 1)$
 $-\Delta = t^9 - t^3 = (t(t^2 - 1))^3$
32. **II II 6** $y^2 = x^3 + tx^2 + t(2t^2 + t + 2)x + t(t^4 + t^3 + t^2 + t + 1)$
 $-\Delta = t^9 + t^6 + t^3 = t^3(t-1)^6$
33. **II II 5 1** $y^2 = x^3 + tx^2i + t(t^2 + t + 2)x + t^5i$
 $-\Delta = 2t^9 + t^8 + t^7 + 2t^5 + 2t^4 + t^3 = -t^3(t-1)^5(t+1)$
34. **II II 4 2** $y^2 = x^3 + tx^2i + t(2t^2 + t + 2)x + t(t^4i + t^2i + i)$
 $-\Delta = t^9 + t^8 + 2t^7 + t^6 + 2t^5 + t^4 + t^3 = t^3(t-1)^4(t+1)^2$
35. **II II 4 1 2** $y^2 = x^3 + tx^2 + t(2t^2 + t + 2)x + t(2t^4 + t^2 + 2)$
 $-\Delta = t^9 + 2t^8 + t^7 + t^6 + t^5 + 2t^4 + t^3 = t^3(t-1)^4(t - \frac{1}{t})(t + \frac{1}{t})$
36. **II II 3 2** $y^2 = x^3 + tx^2 + t(t^2 + t + 2)x + t(t^4 + 2t^3 + t^2 + 1)$
 $-\Delta = 2t^9 + t^3 = -t^3(t-1)^3(t+1)^3$
37. **II II 3 2 1** $y^2 = x^3 + tx^2 + t(t^2 + t + 2)x + t((2-i)t^4 - t^3i + (1+i)t^2 - ti + 2i)$
 $-\Delta = 2t^9 + (2+i)t^8 + (2+i)t^7 - it^6 + (1+i)t^5 + (1+i)t^4 + t^3 = -t^3(t - \frac{1}{t})^3(t-1)^2(t + \frac{1}{t})$
38. **II II 3 1 3** $y^2 = x^3 + tx^2 + t(t^2 + 1)x + t(t^4 + 2t^3 + 2)$
 $-\Delta = 2t^9 + t^7 + 2t^6 + 2t^4 + 2t^3 = -t^3(t-1)^3(t^3 + 2t + 2)$
39. **II II 2 3** $y^2 = x^3 + tx^2i + t(2t^2 + t + 2)x + t(t^4i - t^3i - ti + i)$
 $-\Delta = t^9 + t^8 + 2t^6 + t^4 + t^3 = t^3(t-1)^2(t - \frac{1}{t})^2(t + \frac{1}{t})^2$
40. **II II 2 2 1 2** $y^2 = x^3 + tx^2 + t(2t^2 + t + 2)x + t(t^4 + 2t^3 + 2t^2 + 2t + 1)$
 $-\Delta = t^9 + 2t^7 + 2t^5 + t^3 = t^3(t-1)^2(t+1)^2(t - \frac{1}{t})(t + \frac{1}{t})$
41. **II II 2 1 4** $y^2 = x^3 + tx^2 + t(2t^2 + t + 2)x + t(t^4 + t^3 + 2t^2 + 2t + 2)$
 $-\Delta = t^9 + 2t^5 + 2t^4 + t^3 = t^3(t-1)^2(t^4 - t^3 + t + 1)$
42. **II II 1 6** $y^2 = x^3 + tx^2 + t(t^2 + t + 2)x + t(t^4 + t^3 + t^2 + 2t + 1)$
 $-\Delta = 2t^9 + 2t^7 + 2t^5 + t^3 = -t^3(t-1)^2(t+1)(t^4 + 1)$

4.2.2 Case 1B

$$\delta = 4, \quad r = 0 \quad \text{Type II}$$

$$\begin{aligned} y^2 &= x^3 + tc_1x^2 + t^2c_2x + tc_5, \quad t \nmid c_5, t \nmid c_1 \\ -\Delta &= t^2c_1^2(t^2c_1c_5 - t^4c_2^2) + t^6c_2^3 \end{aligned} \quad (4.2.2)$$

1. **II 8** does not exist, F
2. **II 7 1** $y^2 = x^3 + t(t+1)x^2 + t^2x + t(2+t)$
 $-\Delta = 2t^5 + t^4 = -t^4(t-1)$
3. **II 6 2** does not exist, F
4. **II 6 1²** $y^2 = x^3 + t(t+1)x^2 + 2t^2x + t(2+t)$
 $-\Delta = 2t^6 + 2t^5 + t^4 = t^4(-t^2 - t + 1)$
5. **II 5 3** does not exist, F
6. **II 5 2 1** $y^2 = x^3 + t(t+i)x^2 + (1-i)t^2x + t(2+ti)$
 $-\Delta = 2t^7 + (-1+i)t^6 + 2t^5 - it^4 = t^4(-i - t - (1-i)t^2 - t^3)$
7. **II 5 1³** $y^2 = x^3 + t(t+1)x^2 + t^2x + t(t+1)$
 $-\Delta = t^7 + 2t^5 + 2t^4 = t^4(2 - t + t^3)$
8. **II 4²** $y^2 = x^3 + t(t+1)x^2 + t^2x + t(1+2t)$
 $-\Delta = 2t^8 + t^7 + t^5 + 2t^4 = -t^4(t-1)^4$
9. **II 4 3 1** $y^2 = x^3 + t(t+1)x^2 + t^2x + t((2-i)t + 1 + i)$
 $-\Delta = (-1+i)t^8 - (2+i)t^7 + (-2+i)t^5 - (1+i)t^4 = -(1-i)t^4(t-1) \left(t - \frac{1}{i}\right)^3$
10. **II 4 2²** $y^2 = x^3 + t(t+i)x^2 + t^2x - it$
 $-\Delta = t^8 + t^6 + t^4 = t^4(t-1)^2(t+1)^2$
11. **II 4 2 1²** $y^2 = x^3 + t(t+1)x^2 + t^2xi + t(ti+1+i)$
 $-\Delta = (2+2i)t^8 - it^7 + (2+i)t^6 - it^5 + (2+2i)t^4 = t^4(t-1)^2((2+2i)t^2 + t + (2+2i))$
12. **II 4 1⁴** $y^2 = x^3 + t(t+1)x^2 + 2t^2x + t$
 $-\Delta = t^8 + t^7 + 2t^6 + 2t^4 = -t^4(2 - t + t^2 - t^3)(t-1)$
13. **II 3² 2** does not exist, F
14. **II 3² 1²** $y^2 = x^3 + t(t+1)x^2 + t^2xi + t(2 + (1-i)t^2 + t)$
 $-\Delta = (-1+i)t^9 + t^8 + 2t^7 + (1-i)t^6 + 2t^5 + t^4 = -t^4(t-1)^3((1-i)t^2 - t + 1)$
15. **II 3 2² 1** $y^2 = x^3 + t(t+i)x^2 + 2t^2x + t(1 - (1+i)t^2 + (1+i)t)$
 $-\Delta = (1+i)t^9 + 2it^8 - (1+i)t^7 + (1-i)t^6 + (2+i)t^5 + t^4i = it^4(t-1)^2 \left(t - \frac{1}{1-i}\right)^2 ((1-i)t + i)$
16. **II 3 2 1³** $y^2 = x^3 + t(t+1)x^2 + t^2x + t(2+t^2)$
 $-\Delta = 2t^9 + t^8 + 2t^6 + t^4 = t^4(t-1)^2(1-t-t^2-t^3)$

17. **II 3 1⁵** $y^2 = x^3 + t(t+i)x^2 + t^2x + t(1 + (2-i)t^2)$
 $-\Delta = (-2+i)t^9 + t^8 + (2-i)t^7 - (1+i)t^6 + t^4i = -t^4(t-1)(t+1)(i - t^2 + (2-i)t^3)$
18. **II 2⁴** $y^2 = x^3 + t(t+\epsilon)x^2 + t^2x + t(-it^3 + t^2(\epsilon + \epsilon i) + (1-i)t + 2i\epsilon)$
 $-\Delta = t^{10}i + (2\epsilon - i\epsilon)t^9 + it^8 + (i\epsilon + \epsilon)t^7 + 2it^6 + (2\epsilon - i\epsilon)t^5 + 2it^4 = it^4 \left(t - \frac{1}{i}\right)^2 \left(t + \frac{1}{i}\right)^2 \left(t - \frac{\epsilon}{1+i}\right)^2$, where $\epsilon^4 = -1$
19. **II 2³ 1²** $y^2 = x^3 + t(t+1)x^2 + t^2x + t(2t^3 + (1+i)t^2 + ti + i)$
 $-\Delta = t^{10} + (-1+2i)t^9 - (2+i)t^8 - it^7 + (2+2i)t^6 + 2it^5 - it^4 = -t^4(t-1)^2 \left(t - \frac{1}{i}\right)^2 (-t^2 - t - i)$
20. **II 2² 1⁴** $y^2 = x^3 + t(t+1)x^2 + 2t^2x + t(t^3 + 2t^2 + 1)$
 $-\Delta = 2t^{10} + t^9 + t^8 + 2t^4 = t^4(t-1)^2(2+t-t^3-t^4)$
21. **II 2 1⁶** $y^2 = x^3 + t(t+1)x^2 + t^2x + t(2 + 2t^3 + 2t^2)$
 $-\Delta = t^{10} + t^9 + t^8 + t^7 + t^6 + t^4 = -t^4(t-1) \left(t - \frac{1}{i}\right) \left(t + \frac{1}{i}\right) (1+t+t^2-t^3)$
22. **II 1⁸** $y^2 = x^3 + t(t+1)x^2 + t^2x + t(t^4i - it^3 + (1+i)t^2 + (1-i)t + 1)$
 $-\Delta = -it^{11} + t^{10}i - (1+i)t^9 + (-2+i)t^7 + (2-i)t^6 + (-1+i)t^5 + 2t^4 = -t^4(t-1)(t+i)(t-i)(2 + (1+i)t + t^2 + it^4)$
23. **III 5** $y^2 = x^3 + tx^2 + t^3x + t(2t^4 + t^3 + 2t^2 + 2t^2 + 2)$
 $-\Delta = 2t^9 + 2t^8 + 2t^7 + t^6 + t^5 + t^4 = -t^4(t-1)^5$
24. **III 4 1** $y^2 = x^3 + tx^2 + t^2(2t+1)x + t(t^4 + 2t^3 + t^2 + 2)$
 $-\Delta = t^9 + 2t^7 + 2t^6 + t^4 = t^4(t-1)^4(t+1)$
25. **III 3 2** $y^2 = x^3 + tx^2i + t^3x + t(t^4i + t^3i - t^2i + ti + 2i)$
 $-\Delta = 2t^9 + t^8 + 2t^7 + t^6 + 2t^5 + t^4 = -t^4(t-1)^3(t+1)^2$
26. **III 3 1²** $y^2 = x^3 + tx^2 + t^3x + t(t^4 + t^3 + 2t^2 + 2)$
 $-\Delta = 2t^9 + 2t^7 + t^6 + t^4 = -t^4(t-1)^3 \left(t - \frac{1}{i}\right) \left(t + \frac{1}{i}\right)$
27. **III 2² 1** $y^2 = x^3 + tx^2i + t^3x + t(t^4i - t^3 + t^2i + ti + 2i)$
 $-\Delta = 2t^9 + t^8 + t^7 + 2t^6 + 2t^5 + t^4 = -t^4 \left(t - \frac{1}{i}\right)^2 \left(t + \frac{1}{i}\right)^2 (t-1)$
28. **III 2 1³** $y^2 = x^3 + tx^2 + t^2(t+1)x + t(2t^4 + t^3 + 2)$
 $-\Delta = 2t^9 + 2t^8 + t^7 + t^5 = t^4(t-1)^2(2t^3 + 2t + 1)$
29. **III 1⁵** $y^2 = x^3 + tx^2 + t^3x + t(t^4 + t^3 + t^2 + t + 2)$
 $-\Delta = 2t^9 + 2t^7 + 2t^6 + 2t^5 + t^4 = -t^4(t-1)(t+1)(t^3 - t + 1)$
30. **III 5** $y^2 = x^3 + tx^2i + t^3x + t(t^3i - t^2i - ti + 2i)$
 $-\Delta = 2t^9 + 2t^8 + 2t^7 + t^6 + t^5 + t^4 = -t^4(t-1)^5$
31. **III 4 1** $y^2 = x^3 + t\epsilon x^2 - t^3xi + \frac{t}{\epsilon}((2+2i)t^3 + (2+i)t^2 + t + 2 + 2i)$
 $-\Delta = -t^9i + (1-i)t^8 + t^7 + (3-i)t^6 + (1-i)t^5 + t^4 = -t^4i(t+1)^4 \left(t - \frac{1}{i}\right)$
32. **III 3 2** $y^2 = x^3 + tx^2 + t^3x + t(t^3 + 2t^2 + t + 2)$
 $-\Delta = 2t^9 + t^8 + 2t^7 + t^6 + 2t^5 + t^4 = -t^4(t-1)^3(t+1)^2$

33. **II III 3 1²** $y^2 = x^3 + tx^2 + t^2(2t+1)x + t(2t^3 + t^2 + t + 2)$
 $-\Delta = t^9 + t^8 + 2t^7 + 2t^6 + 2t^5 + t^4 = -t^4(t-1)^3(2t^2 + 2t + 1)$
34. **II III 2² 1** $y^2 = x^3 + tx^2 + t^3x + t(2t^3 + t^2 + t + 2)$
 $-\Delta = 2t^9 + t^8 + t^7 + 2t^6 + 2t^5 + t^4 = -t^4(t-1)\left(t - \frac{1}{t}\right)^2\left(t + \frac{1}{t}\right)^2$
35. **II III 2 1³** $y^2 = x^3 + tx^2 + t^2(2t+1)x + t(t^3 + 2t^2 + 2t + 1)$
 $-\Delta = t^9 + t^8 + t^6 + t^5 + 2t^4 = t^4(t-1)^2(t^3 + 2t + 2)$
36. **II III 1⁵** $y^2 = x^3 + tx^2 + t^2(t+1)x + t(t^3 + 2t^2 + 2)$
 $-\Delta = 2t^9 + t^8 + t^7 + t^6 + t^4 = -t^4(t-1)(t+1)(t^3 + 2t^2 + 1)$

4.2.3 Case 1C

$$\delta = 6, \quad r = 0 \quad \text{Type II}$$

$$y^2 = x^3 + t^2c_0x^2 + t^2c_2x + tc_5, \quad t \nmid c_5, t \nmid c_2 \quad (4.2.3)$$

$$-\Delta = t^4c_0^2(t^3c_0c_5 - t^4c_2^2) + t^6c_2^3$$

1. **II 6** does not exist, G
2. **II 5 1** $y^2 = x^3 + t^2x^2 + t^2(t^3 + t + 1)x + t(2t^5 + t^3 + t^2 + 2)$
 $-\Delta = 2t^{12} + t^{11} + t^{10} + 2t^8 + 2t^7 + t^6 = -t^6(t-1)^5(t+1)$
3. **II 4 2** $y^2 = x^3 + t^2x^2 + t^2(t^2 + t + 2)x + t(2t^5 + t^4 + 2 + 2t^2 + 2t)$
 $-\Delta = t^4(t^3(2t^5 + t^4 + 2t^2 + 2t + 2) + 2t^4(t^2 + t + 2)^2 + t^6(t^2 + t + 2)^3) = -t^6(t-1)^4(t+1)^2$
4. **II 4 1²** $y^2 = x^3 + t^2x^2 + t^2(t^2 + t + 2)x + t(2t^5 + t^3 + 2t^2 + 1)$
 $-\Delta = 2t^{12} + t^{11} + 2t^{10} + 2t^9 + 2t^8 + t^7 + 2t^6 = -t^6(t-1)^4(t-i)(t+i)$
5. **II 3²** does not exist, G
6. **II 3 2 1** $y^2 = x^3 + t^2x^2 + (2t^4 + t^3 + t^2)x + it^5 + t^4 + 2t^3 + 2(1+i)t^2 + t$
 $-\Delta = t^{12} + (2+i)t^{11} + 2t^{10} + t^9 + (1+2i)t^8 + t^7 + t^6 = -t^6(t-1)^3(-t^3 + (1-i)t^2 + t + 1)$
7. **II 3 1³** $y^2 = x^3 + t^2x^2 + t^2(1 + t^2 + t)x + t(t^5 + 2t^4 + 2t^3 + 2t^2 + t + 1)$
 $-\Delta = t^{12} + 2t^{10} + t^9 + t^7 + t^6 = -t^6(t-1)^3(2t^3 + t + 1)$
8. **II 2³** $y^2 = x^3 + 2t^2x^2 + (t^4 + t^3 + \phi_1t^2)x + t^2\phi_1^3 + 2t^5 + (2 + \phi_1)t^4 + (2 + \phi_1)t^3 + 2\phi_1^2t^2 + t$
 $-\Delta = 2t^{12}\phi_1^3 + 2t^{11} + 2t^9 + 2t^7 + t^6\phi_1^3 = -t^6(t-1)^2(t+1)^2(2t^2\phi_1^3 + 2t + \phi_1^3),$
where $\phi_1^6 = -1$
9. **II 2² 1²** $y^2 = x^3 + t^2x^2 + (t^4 + t^3 + 2t^2)x + t^6 + 2t^4 + t^3 + t^2 + t$
 $-\Delta = t^{12} + t^{11} + t^9 + t^7 + 2t^6 = -t^6(t-1)^2(t+1)^2(2t^2 + 2t + 1)$
10. **II 2 1⁴** $y^2 = x^3 + t^2x^2 + (t^4 + t^3 + t^2)x + t^6 + t^5 + t^4 + t^3 + t^2 + t$
 $-\Delta = t^{12} + 2t^{11} + t^{10} + t^7 + t^6 = -t^6(t-1)^2(t+1)(2t^3 + t + 2)$

11. **II 1⁶** $y^2 = x^3 + t^2x^2 + t^2(t^2 + t + 1)x + t(2t^5 + t^3 + 2t^2 + 1)$
 $-\Delta = t^4(t^3(2t^5 + t^3 + 2t^2 + 1) + 2t^4(t^2 + t + 1)^2) + t^6(t^2 + t + 1)^3 =$
 $-t^6(t - 1)(t + 1)(t^4 + 2t^3 + t + 1)$
12. **II II** $y^2 = x^3 + t^2x + t(t^4 + t^3 + t^2 + t + 1)$
 $-\Delta = 2t^6$
13. **II III** $y^2 = x^3 + t^2(2t + 1)x + t(t^4 + t^3 + t^2 + 2t + 1)$
 $-\Delta = t^9 + 2t^6 = 2t^6(2t + 1)^3$
14. **II III III** $y^2 = x^3 + t^2(2t + 1)x + t(t^4 + t^3 + 2t^2 + t + 1)$
 $-\Delta = t^9 + 2t^6 = 2t^6(2t + 1)^3$
15. **II III III** $y^2 = x^3 + t^2(2t + 1)x + t(t^2 + t + 1)$
 $-\Delta = t^9 + 2t^6 = 2t^6(2t + 1)^3$

4.2.4 Case 1D

$\delta = 7, \quad r = 0 \quad \text{Type II}$

$$y^2 = x^3 + t^2c_0x^2 + t^3c_1x + tc_5, \quad t \nmid c_5, t \nmid c_0 \quad (4.2.4)$$

$$-\Delta = t^4c_0^2(t^3c_0c_5 - t^6c_1^2) + t^9c_1^3$$

1. **II 5** $y^2 = x^3 + t^2x^2 + t^2(t + 1)x + t(2t^5 + t^4 + t + 1)$
 $-\Delta = t^4(t^3(2t^5 + t^4 + t + 1) + 2t^6(t + 1)^2) + t^9(t + 1)^3 = -t^7(t - 1)^5$
2. **II 4 1** $y^2 = x^3 + t^2x^2 + t^2(t + 1)x + t(2t^5 + t^4 + 2t^3 + 2)$
 $-\Delta = 2t^{12} + t^{10} + t^9 + 2t^7 = -t^7(t + 1)(t - 1)^4$
3. **II 3 2** $y^2 = x^3 + t^2x^2 + t^2(t + 1)x + t(2t^5 + 2t + 1)$
 $-\Delta = 2t^{12} + t^{11} + 2t^{10} + t^9 + 2t^8 + t^7 = -t^7(t - 1)^3(t + 1)^2$
4. **II 3 1²** $y^2 = x^3 + 2t^2x^2 + (t^3 + t^2)x + t^6 + t^4 + t^2 + t$
 $-\Delta = 2t^{12} + t^{11} + t^{10} + t^9 + 2t^8 + 2t^7 = -t^7(t - 1)^3(t^2 + 2t + 2)$
5. **II 2 1³** $y^2 = x^3 + 2t^2x^2 + t^2(t + 1)x + t(2t^5 + t^2 + 1)$
 $-\Delta = t^4(2t^3(2t^5 + t^2 + 1) + 2t^6(t + 1)^2) + t^9(t + 1)^3 = -t^7(t - 1)^2(t + 1)(2t^2 + t + 1)$
6. **II 2² 1** $y^2 = x^3 + t^2x^2 + t^2(t + 1)x + t(2t^5 + 2t^3 + t^2 + 2t + 1)$
 $-\Delta = 2t^{12} + t^{11} + t^{10} + 2t^9 + 2t^8 + t^7 = -t^7(t - i)^2(t + i)^2(t - 1)$
7. **II 1⁵** $y^2 = x^3 + t^2x^2 + t^2(t + 1)x + t(t^5 + 2t^4 + 2t^3 + t^2 + t + 1)$
 $-\Delta = t^4(t^3(t^5 + 2t^4 + 2t^3 + t^2 + t + 1) + 2t^6(t + 1)^2) + t^9(t + 1)^3 =$
 $-t^7(t - 1)(t + 1)(2t^3 + t + 1)$
8. **II IV** does not exist, E
9. **II II 1** does not exist, E
10. **II II 2** does not exist, E

11. **IIII1²** does not exist, E
12. **IIIII2** does not exist, E
13. **IIIII1²** does not exist, E

4.2.5 Case 1E

$$\delta = 9, \quad r = 0 \quad \text{Type II}$$

$$\begin{aligned} y^2 &= x^3 + t^3 c_1 x + t c_5, \quad t \nmid c_5, t \nmid c_1 \\ -\Delta &= t^9 c_1^3 \end{aligned} \quad (4.2.5)$$

1. **II3** does not exist, E
2. **II21** does not exist, E
3. **II1³** does not exist, E
4. **IIII** $y^2 = x^3 + t^3(2t+1)x + t(t^3+t+1)$
 $-\Delta = t^9(2t+1)^3$
5. **IIIII** $y^2 = x^3 + t^3(2t+1)x + t(t^2+t+1)$
 $-\Delta = t^9(2t+1)^3$

4.2.6 Case 1F

$$\delta = 12, \quad r = 0 \quad \text{Type II}$$

$$\begin{aligned} y^2 &= x^3 + t^4 c_0 x + t c_5, \quad t \nmid c_5, t \nmid c_0 \\ -\Delta &= t^{12} c_0^3 \end{aligned} \quad (4.2.6)$$

4.2.7 Case 2

$$\delta = 3, \quad r = 1 \quad \text{Type III}$$

$$\begin{aligned} y^2 &= x^3 + t c_1 x^2 + t c_3 x + t^2 c_4, \quad t \nmid c_3 \\ -\Delta &= t^2 c_1^2 (t^3 c_1 c_4 - t^2 c_3^2) + t^3 c_3^3 \end{aligned} \quad (4.2.7)$$

1. **IIIIIIIIII** $y^2 = x^3 - t(t-1)(t - (\alpha + \alpha^3))x + (\alpha^5 - \alpha^3)(t^2(t-1)^2)$
 $-\Delta = t^9 - t^6(1 + (\alpha^3 - \alpha)) + (\alpha^3 - \alpha)t^3 = t^3(t-1)^3(t - (\alpha + \alpha^3))^3$,
where $\alpha^4 + \alpha^2 = 1$.
2. **IIIIIIIIII** $y^2 = x^3 - t(t^2 - 1)x + t^2(t-1)^2$
 $-\Delta = t^9 + t^3 = t^3(t^2 + 1)^3$
3. **IIIIIIIIII** $y^2 = x^3 - t(t^2 + 1)x + t^2(t^2 + 1)^2$
 $-\Delta = t^9 + t^3 = t^3(t^2 + 1)^3$

4. **IIIIIIII** $y^2 = x^3 - t(t^2 + 1)x + t^2(t^2 + 1)(t - 1)$
 $-\Delta = t^9 + t^3 = t^3(t^2 + 1)^3$
5. **IIII6** does not exist, G
6. **IIII51** $y^2 = x^3 + tx^2i + t(t^2 + 2)x + t^2(t^3i - t^2i - ti + i)$
 $-\Delta = 2t^9 + t^8 + t^7 + 2t^5 + 2t^4 + t^3$
7. **IIII42** $y^2 = x^3 + tx^2i + t(t^2 + 1)x + t^2(t^3 + t^2i + (1 - i)t + 2 + 2i)$
 $-\Delta = 2t^9 - (1 - i)t^8 - t^7 + (1 - i)t^6 + (1 - i)t^5 + t^4 + t^3i$
8. **IIII41²** $y^2 = x^3 + tx^2 + t(2t^2 + 1)x + t^2(2t^3 + t^2 + t + 2)$
 $-\Delta = t^9 + 2t^8 + 2t^7 + t^5 + t^4 + 2t^3 = t^3(t - 1)^4(t^2 + 2)$
9. **IIII3²** does not exist, G
10. **IIII321** $y^2 = x^3 + tex^2 + t(2t^2 + 2)x + \frac{t^2}{\epsilon}((1 - i)t^3 + t^2 + it + i)$
 $-\Delta = t^9 + (1 + i)t^8 + (2 + i)t^7 + t^6 + (2 - i)t^5 + (1 - i)t^4 + t^3 = t^3(t - 1)^3(t - i)^2(t + 1)$
11. **IIII31³** $y^2 = x^3 + tx^2 + t(t^2 + 1)x + t^2(t^3 + t^2)$
 $-\Delta = 2t^9 + 2t^7 + 2t^6 + t^4 + 2t^3 = -t^3(t - 1)^3(t^3 + t + 2)$
12. **IIII2³** $y^2 = x^3 + tx^2 + t((1 - i)t^2 + 1 + i)x + t^2(-it^3 + 2t^2 + (1 + i)t + 2)$
 $-\Delta = (-1 - i)t^9 - it^8 + t^7 - it^6 + t^5 + 2it^4 + (-1 + i)t^3 = t^3(t - 1)^2(t + 1)^2(-1 + i - it + (-1 - i)t^2)$
13. **IIII2²1²** $y^2 = x^3 + tx^2 + t(t^2i + 1)x + t^2(t^3 + (1 - i)t^2 - (1 + i)t + 1 - i)$
 $-\Delta = t^9i + t^8 + (2 + i)t^7 + t^6 - (1 + 2i)t^5 + t^4 - t^3 = t^3(t - 1)^2(t + 1)^2(2 + t + it^2)$
14. **IIII21⁴** $y^2 = x^3 + tx^2 + t(t^2 + 1)x + t^2(t^3 + 2 + 2t)$
 $-\Delta = 2t^9 + t^5 + t^4 + 2t^3 = t^3(t - 1)^2(2 + 2t + t^3 + 2t^4)$
15. **IIII1⁶** $y^2 = x^3 + tx^2 + t(2t^2 + 2)x + t^2(2t^3 + t^2 + 2t + 1)$
 $-\Delta = t^9 + 2t^8 + 2t^7 + 2t^5 + t^4 + t^3 = t^3(t - 1)(t + 1)(t - i)(t + i)(t - \frac{1}{1-i})(t - \frac{1}{1+i})$
16. **IIII6** does not exist, G
17. **IIII51** $y^2 = x^3 + tex^2 + t^3xi + tx + \frac{t^4}{\epsilon} + \frac{t^3}{(2+2i)\epsilon} + \frac{2t^3i}{2+2i\epsilon} + \frac{2t^2}{\epsilon}$
 $-\Delta = t^9i - (2 + 2i)t^8 + (1 + 1i)t^7 + (\frac{1}{1+i} + 2 + i)t^6 + (2 + 2i)t^5 + (2 + 2i)t^4 + 2t^3$, where $\epsilon^2 + 1 + i = 0$
18. **IIII42** does not exist, G
19. **IIII41²** $y^2 = x^3 + tex^2 + t(t^2 + 1 + i)x + \frac{t^2}{\epsilon}(-t^2i - (\frac{1}{2} - \frac{1i}{2})t + 1)$
 $-\Delta = 2t^9 - (1 + i)t^8 + (1 - i)t^7 - (1 + i)t^6 - (2 + 2i)t^5 + (2 + i)t^4 + (2 + i)t^3$,
where $\epsilon^2 + 1 + i = 0$
20. **IIII3²** does not exist, G
21. **IIII321** $y^2 = x^3 + tex^2 + t((1 - i)t^2 + 2)x + \frac{t^2}{\epsilon}(-t^2 + (\frac{1}{5} - \frac{2i}{5})t + 1)$
 $-\Delta = (2 + 2i)t^9 + (2 + 2i)t^8 + 2it^7 + t^6 + 2it^5 - (1 - i)t^4 + t^3$

22. **III III 3 1³** $y^2 = x^3 + tx^2 + t(2t^2 + 1)x + t^2(1 + t^2 + t)$
 $-\Delta = t^9 + t^8 + 2t^7 + 2t^5 + t^4 + 2t^3 = -t^3(t-1)^3(2t^3 + 2t^2 + t + 2)$
23. **III III 2³** $y^2 = x^3 + tx^2 + t(2t^2 + t + 2)x + t^2(1 + t^2)$
 $-\Delta = t^9 + t^8 + 2t^6 + t^4 + t^3 = t^3(t-1)^2(t-i)^2(t+i)^2$
24. **III III 2² 1²** $y^2 = x^3 + tx^2 + t(t^2 + 2)x + t^2(t^2 + 2t + 2)$
 $-\Delta = 2t^9 + t^8 + 2t^7 + 2t^6 + t^5 + t^4 + t^3 = -t^3 \left(\frac{t-1}{i}\right)^2 \left(\frac{t+1}{i}\right)^2 \left(t - \frac{1}{1+i}\right) \left(t - \frac{1}{1-i}\right)$
25. **III III 2 1⁴** $y^2 = x^3 + tx^2 + t(2t^2 + 1)x + t^2(t^2i + ti + i)$
 $-\Delta = t^9 + 2t^8 + 2t^7 + t^6 + 2t^5 + 2t^4 + 2t^3 = t^3(t-1)^2(-1 + t^3 + t^4)$
26. **III III 1⁶** $y^2 = x^3 + tx^2 + t(t^2 + t + 1)x + t^2(1 + t^2)$
 $-\Delta = 2t^9 + t^8 + 2t^7 + t^5 + t^4 + 2t^3 = -t^3(t-1)(t^5 + t^3 + t^2 + 2)$
27. **III 9** does not exist, A
28. **III 8 1** does not exist, F
29. **III 7 2** does not exist, F
30. **III 7 1²** $y^2 = x^3 + t(t+1)x^2 + t(2t^2 + 2)x + t^4$
 $-\Delta = 2t^5 + t^4 + t^3 = t^3(1 + t - t^2)$
31. **III 6 3** $y^2 = x^3 + t^2x^2 + t(t^2 + 1)x + t^2(2 + t^2 + 2t)$
 $-\Delta = t^6 + 2t^3 = t^3(t-1)^3$
32. **III 6 2 1** does not exist, F
33. **III 6 1³** $y^2 = x^3 + t(t+1)x^2 + t(t^2 + 1)x + t^2(t^2 + t)$
 $-\Delta = 2t^6 + 2t^5 + t^4 + 2t^3 = t^3(2 + t - t^2 - t^3)$
34. **III 5 4** does not exist, F
35. **III 5 3 1** does not exist, F
36. **III 5 2²** does not exist, C
37. **III 5 2 1²** $y^2 = x^3 + tx^2 + t(-it^3 + 2)x + t((1+i)t^5 + 2t^4 + (1-i)t^3 + t^2i + t)$
 $-\Delta = 2it^{12} + 2t^{10} + (-1-i)t^9 + t^8 + 2t^7 + 2it^6 + 2t^5 + t^4 + t^3$
38. **III 5 1⁴** $y^2 = x^3 + t(t+1)x^2 + t(2t^2 + i)x + t^2(t^2 + 1 + i)$
 $-\Delta = (-1-i)t^7 + (2+i)t^6 - it^5 + 2t^4 + t^3i = -t^3(t-1)(i - (1-i)t - t^2 + (1+i)t^3)$
39. **III 4² 1** does not exist G
40. **III 4 3 2** does not exist, F
41. **III 4 3 1²** $y^2 = x^3 + tx^2 + t((-1-i)t^3 + 1 + 2i)x + t(2t^3 + (1-i)t^2)$
 $-\Delta = (1-i) * t^{12} + 2it^{10} + (2+i)t^6 - 2it^4 - (1+i)t^3$
42. **III 4 2² 1** $y^2 = x^3 + t(t+1)x^2 + t((-1-i)t^2 + 2 + 2i)x + t^2(-it^2 + t + i)$
 $-\Delta = -it^8 + 2t^6 + 2it^4 + (1-i)t^3 = -t^3(t-1)^2 \left(t + \frac{1}{i}\right)^2 (it + (1-i))$

43. **III 4 2 1³** $y^2 = x^3 + t(t+1)x^2 + t(t^2+1)x + t^2(2+t^2+t)$
 $-\Delta = t^8 + 2t^6 + t^4 + 2t^3 = t^3(t-1)^2(2-t-t^2+t^3)$
44. **III 4 1⁵** $y^2 = x^3 + t(t+1)x^2 + t(t^2+1)x + t^2(1+t^2+t)$
 $-\Delta = 2t^8 + 2t^6 + t^5 + t^4 + 2t^3 = t^3(2+t+t^2-t^3-t^5)$
45. **III 3³** $y^2 = x^3 + t^2x^2 + t(t^2+2)x + t^2(1+t^2)$
 $-\Delta = 2t^9 + t^6 + t^3 = -t^3 \left(t - \frac{1}{(1-i)}\right)^3 \left(t - \frac{1}{(1+i)}\right)^3$
46. **III 3² 2 1** does not exist, F
47. **III 3² 1³** $y^2 = x^3 + (t^2 + ti)x^2 + (\phi_0 t + t^3 i)x + 2t^4 + t^3 \phi_0$
 $- = (2i + 2\phi_0)t^9 + (-i\phi_0 + 1)t^8 + (2i + 2\phi_0)t^7 + (2i + 2\phi_0 - i\phi_0)t^6 +$
 $(\phi_0 i + 2)t^5 + (i + \phi_0)t^4 + (2\phi_0 + 2i + \phi_0 i)t^3 = -t^3(t-1)^3((\phi_0 + i)t^3 +$
 $(-1 + i\phi_0)t^2 + (i + \phi_0)t + (-\phi_0 - i + i\phi_0))$, where $\phi_0^2 + \phi_0 + i = 0$
48. **III 3 2³** $y^2 = x^3 + t(t+1)x^2 + 2tx + t^2(2+2t)$
 $-\Delta = t^9 + t^8 + 2t^6 + t^4 + t^3 = t^3(t-1)^2(t-i)^2(t+i)^2$
49. **III 3 2² 1²** $y^2 = x^3 + tx^2 + t((1+2i)t^3 + 1+i)x + t(t^5i + t^4i + t^3 + (1-i)t^2 - it)$
 $-\Delta = ((-1-i)t^{12} + it^{10} + 2it^9 + 2it^8 + (2+i)t^6 + it^5 + 2it^4 + (-1+i)t^3 =$
 $t^3(t-1)^3(t^6 + (1+i)t^4 - it^3 - (1+2i)t^2 + (1+i)t + i)$
50. **III 3 2 1⁴** $y^2 = x^3 + t(t+1)x^2 + t((1+i)t^2+1)x + t^2(-it^2 + (1+i)t + 1+i)$
 $-\Delta = (1+i)t^9 - (2+3i)t^8 + (1+2i)t^7 + (-1+i)t^6 + (4-i)t^5 + t^4 + 2t^3 =$
 $t^3(t-1)^2(2-t-it^2-it^3 + (1+i)t^4)$
51. **III 3 1⁶** $y^2 = x^3 + t^2x^2 + tx + t^2(1+t)$
 $-\Delta = t^9 + t^8 + -t^6 + t^3 = t^3(t^6 + t^5 - t^3 + 1)$
52. **III 2⁴ 1** $y^2 = x^3 + t\epsilon x^2 + t(2t^3 + 2+i)x + \frac{t^2}{\epsilon}((-1+i)t^4 - (1+i)t^3 -$
 $(1+i)t^2 + ti + i)$
 $-\Delta = t^{12} + (2-i)t^{10} + t^9 + 4it^8 + (2+4i)t^7 + (2+4i)t^6 + (2+4i)t^5 - (2+4i)t^4 + (4+4i)t^3 = t^3(t - (-1-i))^2(t-1)^2(t+1)^2(t+i)^2(t - (-1+i))$,
where $\epsilon^2 = 2 - i$
53. **III 2³ 1³** $y^2 = x^3 + t^2x^2 + tx + t^2(2+2t^2+2t)$
 $-\Delta = t^{10} + t^9 + t^8 + t^6 + 2t^3 = t^3(t-1)^2(t+1)^2(2+t^2+t^3)$
54. **III 2² 1⁵** $y^2 = x^3 + t^2x^2 + t(t^2+2)x + t^2(2+2t^2+2t)$
 $-\Delta = 2t^{10} + 2t^8 + t^6 + t^3 = t^3(t-1)^2(1-t-t^3+t^4-t^5)$
55. **III 2 1⁷** $y^2 = x^3 + (t+t^2)x^2 + 2tx + t^4 + 2t^3$
 $-\Delta = 2t^{10} + t^9 + 2t^7 + 2t^6 + 2t^5 + t^4 + t^3 = t^3(t-i)(t+i)(1+t+t^2+t^3+t^4-t^5)$
56. **III 1⁹** $y^2 = x^3 + t(t+1)x^2 + 2tx + t^2(t^3+2)$
 $-\Delta = 2t^{11} + t^6 + t^4 + t^3 = t^3(t-i)(t+i)(1+t-t^2+t^4-t^6)$

4.2.8 Case 3A

$$\delta = 5, \quad r = 2 \quad \text{Type IV}$$

$$\begin{aligned} y^2 &= x^3 + tc_1x^2 + t^2c_2x + t^2c_4, \quad t \nmid c_4, t \nmid c_1 \\ -\Delta &= t^2c_1^2(t^3c_1c_4 - t^4c_2^2) + t^6c_2^3 \end{aligned} \quad (4.2.8)$$

1. **IV 7** does not exist, C
2. **IV 6 1** does not exist, F
3. **IV 5 2** does not exist, G
4. **IV 5 1²** $y^2 = x^3 + t(t+1)x^2 + 2t^2x + t^2$
 $-\Delta = 2t^7 + 2t^5 = t^5(2t^2 + 2)^2$
5. **IV 4 3** does not exist, F
6. **IV 4 2 1** $y^2 = x^3 + tx^2i + t^2(x^2 + x + 2xi + 2)$
 $-\Delta = (1+i)t^8 + t^7 + t^6 - it^5$
7. **IV 4 1³** $y^2 = x^3 + t(t+1)x^2 + 2t^2xi - t^2(2+i)$
 $-\Delta = -(2+2i)t^8 + t^7 - (1+2i)t^6 - (1+2i)t^5$
8. **IV 3² 1** $y^2 = x^3 + tx^2i + t^2(x^2 + 2i) - t^3i$
 $-\Delta = t^9i + t^8i + t^6 + t^5$
9. **IV 3 2²** $y^2 = -\frac{1}{2}t^3i + (2+2i)(-2xi + x - x^2i + 2 + 2i + x^2)t^2 + (2+2i)(x^2i + x^2)t + (2+2i)(x^3 - x^3i)$
 $-\Delta = -it^9 - (1-i)t^8 + t^7i - it^6 + t^5$
10. **IV 3 2 1²** $y^2 = x^3 + tx^2 + t^2(x^2 + 2x + 1) + 2t^3$
 $-\Delta = t^9 + 2t^7 + t^6 + 2t^5$
11. **IV 3 1⁴** $y^2 = x^3 + t(t+1)x^2 + 2t^2x + t^2(t+1)$
 $-\Delta = 2t^9 + 2t^7 + 2t^6 + 2t^5$
12. **IV 2³ 1** $y^2 = x^3 + t(t+1)x^2 + 2t^2x + t^2((2+i)t^2 + (2+2i)t + 2+2i)$
 $-\Delta = (1+2i)t^{10} + (1+i)t^9 + (2+i)t^8 + (2+i)t^7 + (2-i)t^6 + (1-i)t^5$
13. **IV 2² 1³** $y^2 = x^3 + t(t+1)x^2 + 2t^2x + t^2(t^3\gamma_3 + 2t^2 + t)$
 $-\Delta = t^{10} + 2t^9 + t^8 + 2t^6$
14. **IV 2 1⁵** $y^2 = x^3 + t(t+1)x^2 + t^2x + t^2(t^2 + 2t + 1)$
 $-\Delta = 2t^{10} + t^9 + t^7 + 2t^5$
15. **IV 1⁷** $y^2 = x^3 + t(t+1)x^2 + t^2x + t^2(2t^3 + t^2 + 1)$
 $-\Delta = t^{11} + 2t^{10} + t^8 + t^7 + 2t^6 + 2t^5$
16. **IV IV 2** $y^2 = x^3 + tx^2 + 2t^2x + t^2(2t^2 + 2t + 2)$
 $-\Delta = t^7 + t^6 + t^5 = t^5(t-1)^2$

17. **IV IV 1²** $y^2 = x^3 + tx^2 + 2t^2x + t^2(2 + t^2)$
 $-\Delta = 2t^7 + t^5 = 2t^5(t-1)(t+1)$
18. **IV II 3** $y^2 = x^3 + tx^2 + 2t^2x + t^2(t^3 + 2)$
 $-\Delta = 2t^8 + t^5 = t^5(t-1)^2$
19. **IV II 2 1** $y^2 = x^3 + tx^2 + 2t^3\phi_1x + t^2(2t^3 + t^2 + t + 2)$
 $-\Delta = t^8 + 2t^7 + 2t^6 + t^5$
20. **IV II 1³** $y^2 = x^3 + tx^2 + t^2(2t^3 + t + 2)$
 $-\Delta = t^8 + 2t^6 + t^5$
21. **IV III 4** $y^2 = x^3 + tx^2i + t^3x + t^2(ti + 2i)$
 $-\Delta = t^9 + 2t^8 + 2t^6 + t^5 = t^5(t-1)^4$
22. **IV III 3 1** $y^2 = x^3 + tx^2i + 2t^3x + t^2(2ti + 2i)$
 $-\Delta = 2t^9 + 2t^8 + t^6 + t^5 = -t^5(t-1)^3(t+1)$
23. **IV III 2²** $y^2 = x^3 + 2tx^2 + 2t^2(1+i)t+1)x + t^2((2+2i)t^2 + 2t + 1 + 2i)$
 $-\Delta = -(1-i)t^9 + 2it^8 - (2+2i)t^7 + 2t^6 - (2+i)t^5$
24. **IV III 2 1²** $y^2 = x^3 + tx^2 + 2t^2(2t+1)x + t^2(2t^2 + 2t + 2)$
 $-\Delta = t^9 + t^8 + 2t^7 + t^6 + t^5 = t^5(t-1)^2(t-i)(t+i)$
25. **IV III 1⁴** $y^2 = x^3 + tx^2 + 2t^2(t+2)x + t^2(t^2 + t + 2)$
 $-\Delta = 2t^9 + t^8 + t^6 + t^5 = t^5(2t^4 + t^3 + t + 1)$
26. **IV II 4** $y^2 = x^3 + tx^2 + 2t^2(2t+1)x + t^2(2t^3 + t^2 + t + 2)$
 $-\Delta = t^9 + 2t^8 + 2t^6 + t^5 = t^5(t-1)^4$
27. **IV II 3 1** $y^2 = x^3 + tx^2 + 2t^2(t+1)x + t^2(2t^3 + 2t^2 + 2t + 2)$
 $-\Delta = 2t^9 + 2t^8 + t^6 + t^5 = t^5(t-1)^3(t+1)$
28. **IV II 2²** $y^2 = x^3 + tx^2 + 2t^2(2t+1)x + t^2(t^3 + 2)$
 $-\Delta = t^9 + t^7 + t^5 = t^5(t-1)^2(t+1)^2$
29. **IV II 2 1²** $y^2 = x^3 + tx^2i + t^3x + t^2(t^3i + t^2i + 2ti + 2i)$
 $-\Delta = t^9 + t^8 + 2t^7 + t^6 + t^5 = t^5(t-1)^2(t-i)(t+i)$
30. **IV II 1⁴** $y^2 = x^3 + tx^2 + 2t^2(t+1)x + t^2(2t^3 + 2t^2 + t + 2)$
 $-\Delta = 2t^9 + 2t^8 + 2t^6 + t^5 = t^5(2t^4 + 2t^3 + 2t + 1)$

4.2.9 Case 3B

$$\delta = 6, \quad r = 2 \quad \text{Type IV}$$

$$\begin{aligned} y^2 &= x^3 + t^2c_0x^2 + t^2c_2x + t^2c_4, \quad t \nmid c_4, t \nmid c_2 \\ -\Delta &= t^4c_0^2(t^4c_0c_4 - t^4c_2^2) + t^6c_2^3 \end{aligned} \quad (4.2.9)$$

1. **IV 6** $y^2 = x^3 + t^2x^2 + (t^4 + t^3 + 2t^2)x + 2t^6 + 2t^5 + 2t^3 + t^2$
 $-\Delta = 2t^{12} + 2t^9 + 2t^6 = 2t^6(t^6 + t^3 + 1)$
2. **IV 5 1** does not exist, G
3. **IV 4 2** does not exist, G
4. **IV 4 1²** $y^2 = x^3 + t^2x^2 + (t^4 + t^3 + t^2)x + 2t^6 + t^5 + t^3$
 $-\Delta = 2t^{12} + 2t^{11} + 2t^8 + t^6 = t^6((t-1)^4(t-(-1-i))(t-(-1+i)))$
5. **IV 3²** does not exist, G
6. **IV 3 2 1** does not exist, G
7. **IV 3 1³** $y^2 = x^3 + 2t^2x^2 + (t^3 + t^2)x + 2t^6 + t^5 + 2t^4 + t^3 + t^2$
 $-\Delta = 2t^{12} + 2t^{11} + t^9 + t^8 + t^6 = -t^6(t^3 + 2)(2t^3 + t^2 + 1)$
8. **IV 2³** does not exist, G
9. **IV 2² 1²** $y^2 = x^3 + t^2x^2 + (t^3 + 2t^2)x + 2t^6 + 2t^4 + 2t^2$
 $-\Delta = 2t^{12} + t^{10} + t^8 + 2t^6 = t^6(1 + t^2)(2t^4 + 2t^2 + 2)$
10. **IV 2 1⁴** $y^2 = x^3 + t^2x^2 + t^2(t^2 + t + 1)x + t^2(2t^4 + t^2 + 2t + 1)$
 $-\Delta = 2t^6 + t^5 + t^4 + t^3 + 1 = -t^6(t-1)^2(t-i)(t-(1-i))(t-(1+i))$
11. **IV 1⁶** $y^2 = x^3 + t^2x^2i + t^2(t^2 + t + 1)x + t^2(3it^4 - 2it^3 - 2it^2 + i)$
 $-\Delta = 2t^{12} + t^{10} + 2t^8 + t^6 = -t^6(t-1)(t+1)(t^4 + 1)$
12. **IV II** $y^2 = x^3 + t^2x + t^2(t^3 + 1 + t)$
 $-\Delta = 2t^6$
13. **IV III II** $y^2 = x^3 + t^2(2t + 1)x + t^2(t^3 + t + 1)$
 $-\Delta = t^9 - t^6 = 2t^6(2t + 1)^3$
14. **IV III II** $y^2 = x^3 + t^2(2t + 1)x + t^2(2t + 1)$
 $-\Delta = t^9 - t^6 = 2t^6(2t + 1)^3$
15. **IV III III** $y^2 = x^3 + t^2(2t + 1)x$
 $-\Delta = t^9 - t^6 = 2t^6(2t + 1)^3$
16. **IV IV** $y^2 = x^3 + 2t^2x + t^2(t^2 + t + 1)$
 $-\Delta = t^6$

4.2.10 Case 3C

$$\delta = 8, \quad r = 2 \quad \text{Type IV}$$

$$y^2 = x^3 + t^2c_0x^2 + t^3c_1x + t^2c_4, \quad t \nmid c_4, t \nmid c_0 \quad (4.2.10)$$

$$-\Delta = t^4c_0^2(t^4c_0c_4 - t^6c_1^2) + t^9c_1^3$$

1. **IV 4** $y^2 = x^3 + 2t^2x^2 + t^2(t^4 + 2t^3 + 2t + 1)$
 $\Delta = t^7(t^5 + 2t^4 + 2t^2 + t) = t^{12} + 2t^{11} + 2t^9 + t^8$

2. **IV 3 1** $y^2 = x^3 + t^2x^2 + t^3x + t^2(2t^4 + 2t^3 + t^2 + 1)$
 $\Delta = t^{12} + t^{11} + 2t^8 + 2t^9$
3. **IV 2²** $y^2 = x^3 + t^2x^2 + t^3(t+2)x + t^2(2t^4 + t^3 + 2 + 1)$
 $\Delta = t^{12} + t^8 + t^{10}$
4. **IV 2 1²** $y^2 = x^3 + t^2x^2 + t^3(t+1)x + t^2(2t^4 + t^3 + 2t^2 + t + 2)$
 $\Delta = 2t^7(2t^5 + t) = t^{12} - t^8$
5. **IV 1⁴** $y^2 = x^3 + t^2x^2 + t^2(2t^4 + 1)$
 $-\Delta = t^8(2t^4 + 2t^3 + t^2 + 2t + 2)$
6. **IV II** does not exist, E
7. **IV III 1** does not exist, E
8. **IV III 1** does not exist, E

4.2.11 Case 3D

$\delta = 9, \quad r = 2 \quad \text{Type IV}$

$$\begin{aligned} y^2 &= x^3 + t^3c_1x + t^2c_4, \quad t \nmid c_4, t \nmid c_1 \\ -\Delta &= t^9c_1^3 \end{aligned} \tag{4.2.11}$$

1. **IV 3** does not exist, E
2. **IV 2 1** does not exist, E
3. **IV 1³** does not exist, E
4. **IV II** $y^2 = x^3 + t^3(2t+1)x + t^2$
 $-\Delta = t^9(2t+1)^3$
5. **IV III** $y^2 = x^3 + t^3(2t+1)x + t^2(t^2 + t + 1)$
 $-\Delta = t^9(2t+1)^3$

4.2.12 Case 3E

$\delta = 12, \quad r = 2 \quad \text{Type IV}$

$$\begin{aligned} y^2 &= x^3 + t^4c_0x + t^2c_4, \quad t \nmid c_4, t \nmid c_0 \\ -\Delta &= t^{12}c_0^3 \end{aligned} \tag{4.2.12}$$

4.2.13 Case 4

In [3] there are two different sub-cases to this case, but both have Weierstrass form and discriminant as listed below, and thus we may treat them together.

$$\delta = 6, \quad r = 4 \quad \text{Type } \mathbf{I}_0^*$$

$$\begin{aligned} y^2 &= x^3 + tx^2 + t^2c_2x + t^3c_3 \\ -\Delta &= t^2c_1^2(t^4c_1c_3 - t^8c_0^2) + t^{12}c_0^2 \end{aligned} \quad (4.2.13)$$

1. $\mathbf{I}_0^* \mathbf{6}$ does not exist, A
2. $\mathbf{I}_0^* \mathbf{5} \mathbf{1}$ does not exist, C
3. $\mathbf{I}_0^* \mathbf{4} \mathbf{2}$ does not exist, C
4. $\mathbf{I}_0^* \mathbf{4} \mathbf{1} \mathbf{2}$ $y^2 = x^3 + t^2x^2 + 2t^2x + t^3(t^3 + t^2)$
 $-\Delta = t^4(t^5(t^3 + t^2) + 2t^4) + 2t^6 = t^6(t^6 + t^5 + 2t^2 + 2)$
5. $\mathbf{I}_0^* \mathbf{3} \mathbf{2}$ does not exist, C
6. $\mathbf{I}_0^* \mathbf{3} \mathbf{2} \mathbf{1}$ does not exist, C
7. $\mathbf{I}_0^* \mathbf{3} \mathbf{1} \mathbf{3}$ $y^2 = x^3 + tx^2 + t^4x + t^3(t^3 + t + 1)$
 $-\Delta = t^2(t^4(t^3 + t + 1) + 2t^8) + t^{12} = t^6(t^3 + 2t + 2)(t^3 + 2)$
8. $\mathbf{I}_0^* \mathbf{2} \mathbf{3}$ $y^2 = \left(\frac{1}{\epsilon} + \frac{i}{\epsilon}\right) t^6 - \frac{it^5}{\epsilon} \left(2x + \frac{1-i}{\epsilon}\right) t^4 - \frac{t^3}{\epsilon} + tx^2\epsilon + x^3$
 $-\Delta = t^{12} + it^{10} + (1-i)t^9 + 2t^8 - (1+i)t^7 - 2it^6 = t^6(t+1+i)^2(t-1)^2(t-i)^2$
9. $\mathbf{I}_0^* \mathbf{2} \mathbf{2} \mathbf{1} \mathbf{2}$ $y^2 = x^3 + t^2x^2i + t^2(t^2 + t + 2)x + t^3(t^2i - ti + i)$
 $-\Delta = 2t^{12} + t^{10} + t^8 + 2t^6 = t^6(2t^6 + t^4 + t^2 + 2)$
10. $\mathbf{I}_0^* \mathbf{2} \mathbf{1} \mathbf{4}$ $y^2 = x^3 + tx^2 + t^2(t^2 + t + 1)x + t^3(2t^2 + 2t + 2)$
 $-\Delta = -t^{12} + t^{10} + t^9 + t^8 + t^6 = t^6(t-1)^2(-t^4 + t^3 + t^2 - t + 1)$
11. $\mathbf{I}_0^* \mathbf{1} \mathbf{6}$ $y^2 = x^3 + tx^2 + t^2(2t^2 + 2t + 1)x + t^3(t^2 + t + 2)$
 $-\Delta = t^{12} + t^{10} + t^8 + t^6 = t^6(t-i)(t+i)(t^4 - 1)$
12. $\mathbf{I}_0^* \mathbf{I}_0^*$ $y^2 = x^3 + 2t^2x + t^3$
 $-\Delta = t^6$
13. $\mathbf{I}_0^* \mathbf{IV}$ $y^2 = x^3 + 2t^2x + t^3(t + 1)$
 $-\Delta = t^6$
14. $\mathbf{I}_0^* \mathbf{II}$ $y^2 = x^3 + 2t^2x + t^3(t^2 + t + 1)$
 $-\Delta = t^6$
15. $\mathbf{I}_0^* \mathbf{III} \mathbf{II}$ $y^2 = x^3 + t^2(2t + 1)x + t^3(t^2 + 2t)$
 $-\Delta = t^9 + 2t^6 = t^6(t^3 + 2)$
16. $\mathbf{I}_0^* \mathbf{III} \mathbf{III}$ $y^2 = x^3 + t^2(2t + 1)x$
 $-\Delta = t^9 + 2t^6 = t^6(t^3 + 2)$

17. $\mathbf{I}_0^* \text{II III}$ $y^2 = x^3 + t^2(2t+1)x + t^3(t+2)$
 $-\Delta = t^9 + 2t^6 = t^6(t^3+2)$
18. $\mathbf{I}_0^* \text{IV 1}$ $y^2 = x^3 + tx^2 + t^2x + t^3(t+2)$
 $-\Delta = -t^7 + t^6 = -t^6(t-1)$
19. $\mathbf{I}_0^* \text{II 2}$ $y^2 = x^3 + tx^2 + t^3(t^2+t+1)$
 $-\Delta = t^8 + t^7 + t^6 = t^6(t-1)^2$
20. $\mathbf{I}_0^* \text{II 1}^2$ $y^2 = x^3 + tx^2 + t^3(t^2-1)$
 $-\Delta = t^8 - t^6 = t^6(t-1)(t+1)$
21. $\mathbf{I}_0^* \text{III 3}$ $y^2 = x^3 + tx^2 + t^6 + t^3$
 $-\Delta = t^9 + t^6 = t^6(t+1)^3$
22. $\mathbf{I}_0^* \text{III 2 1}$ $y^2 = x^3 + tx^2i + t^2(2t+1)x$
 $-\Delta = t^9 + 2t^8 + 2t^7 + t^6 = t^6(t-1)^2(t+1)$
23. $\mathbf{I}_0^* \text{III 1}^3$ $y^2 = x^3 + tx^2 + t^2(t+1)x + 2t^3$
 $-\Delta = 2t^9 + t^8 + 2t^7 + t^6 = -t^6(t-1)(t-\frac{1}{i})(t+\frac{1}{i})$
24. $\mathbf{I}_0^* \text{II 3}$ $y^2 = x^3 + tx^2 + t^2(t+1)x + t^3(t^2+2t+2)$
 $-\Delta = 2t^9 + t^6 = -t^6(t-1)^3$
25. $\mathbf{I}_0^* \text{II 2 1}$ $y^2 = x^3 + tx^2 + t^2(2t+1)x + t^3(2t^2+2t+2)$
 $-\Delta = t^9 + 2t^8 + 2t^7 + t^6 = t^6(t-1)^2(t+1)$
26. $\mathbf{I}_0^* \text{II 1}^3$ $y^2 = x^3 + tx^2i + t^2(t+1)x + t^3(t^2i-ti)$
 $-\Delta = 2t^9 + t^8 + 2t^7 + t^6 = -t^6(t-1)(t-i)(t+i)$

4.2.14 Case 5A

$$\delta = 7, \quad r = 5 \quad \text{Type } \mathbf{I}_1^*$$

$$y^2 = x^3 + tc_1x^2 + t^4c_0x + t^4c_2, \quad t \nmid c_1, t \nmid c_2 \quad (4.2.14)$$

$$-\Delta = t^2c_1^2(t^5c_1c_2 - t^8c_0^2) + t^{12}c_0^3$$

1. $\mathbf{I}_1^* \mathbf{5}$ does not exist, A
2. $\mathbf{I}_1^* \mathbf{4 1}$ does not exist, G
3. $\mathbf{I}_1^* \mathbf{3 2}$ does not exist, C
4. $\mathbf{I}_1^* \mathbf{3 1}^2$ $y^2 = x^3 + tx^2 + 2t^4x + t^4(t^2+1)$
 $-\Delta = t^2(t^5(t^2+1) + 2t^8) + 2t^{12} = t^7(t^2+1)(2t^3+1)$
5. $\mathbf{I}_1^* \mathbf{2 1}^3$ $y^2 = x^3 + t(t+1)x^2 + it^4x + t^4((1+i)t+i)$
 $-\Delta = t^2(t+1)^2(t^5(t+1)((1+i)t+i) + t^8) - t^{12}i = t^7(2+i)(2t^3+it^2+it+t+2+i)(t^2+t+1)$
6. $\mathbf{I}_1^* \mathbf{2}^2 \mathbf{1}$ $y^2 = x^3 + t(t+1)x^2 + t^4\alpha x + t^4((2\alpha^3+2\alpha^2)t+2\alpha^3)$
 $-\Delta = t^2(t+1)^2(t^5(t+1)((2\alpha^3+2\alpha^2)+2\alpha^3)+2t^8\alpha^2)+t^{12}\alpha^3 = t^7\alpha^2(2\alpha t^4+2\alpha t^3+2t^3+2t^5+2\alpha t+2t+2\alpha+\alpha t^5), \text{ where } \alpha^4+\alpha+1=0$

7. $\mathbf{I}_1^* \mathbf{1}^5$ $y^2 = x^3 + t(t+1)x^2 + t^4x + t^4(t^2 + 2t + 1)$
 $-\Delta = t^2(t+1)^2(t^5(t+1)(t^2 + 2t + 1) + 2t^8) + t^{12} = t^7(t^5 + t^2 + 2t + 1)$
8. $\mathbf{I}_1^* \mathbf{II} \mathbf{1}^2$ $y^2 = x^3 + t(t+2)x^2 + t^4(t^2 + 1)$
 $-\Delta = t^7(t+2)^3(t^2 + 1)$
9. $\mathbf{I}_1^* \mathbf{II} \mathbf{2}$ $y^2 = x^3 + t(t+2)x^2 + t^4(2t^2 + t + 2)$
 $-\Delta = t^7(t+2)^3(2t^2 + t + 2)$
10. $\mathbf{I}_1^* \mathbf{II} \mathbf{1}$ $y^2 = x^3 + t(t+2)x^2 + t^4(2t^2 + 1)$
 $-\Delta = t^7(t+2)^3(2t^2 + 1)$
11. $\mathbf{I}_1^* \mathbf{IV}$ $y^2 = x^3 + t(t+2)x^2 + t^4(t^2 + t + 1)$
 $-\Delta = t^7(t+2)^3(t^2 + t + 1)$
12. $\mathbf{I}_1^* \mathbf{III} \mathbf{2}$ does not exist, G

4.2.15 Case 5B

$$\delta = 8, \quad r = 6 \quad \text{Type } \mathbf{I}_2^*$$

$$y^2 = x^3 + tc_1x^2 + t^4c_0x + t^5d_1, \quad t \nmid c_1, t \nmid d_1 \quad (4.2.15)$$

$$-\Delta = t^2c_1^2(t^6c_1d_1 - t^8c_0^2) + t^{12}c_0^3$$

1. $\mathbf{I}_2^* \mathbf{4}$ does not exist, A
2. $\mathbf{I}_2^* \mathbf{3} \mathbf{1}$ does not exist, C
3. $\mathbf{I}_2^* \mathbf{2}^2$ $y^2 = x^3 + tx^2 + 2t^4x + 2t^5$
 $-\Delta = t^{12} + t^{10} + t^8 = t^8(t-1)^2(t+1)^2$
4. $\mathbf{I}_2^* \mathbf{2} \mathbf{1}^2$ $y^2 = x^3 + tx^2 + t^4x + t^5(t+2)$
 $-\Delta = 2t^{12} + t^{10} + 2t^9 + t^8 = t^8(t-1)^2(2t^2 + t + 1)$
5. $\mathbf{I}_2^* \mathbf{1}^4$ $y^2 = x^3 + tx^2 + t^4x + t^5(2t+1)$
 $-\Delta = 2t^{12} + t^{10} + t^9 + 2t^8 = t^8(t-1)(2t^3 + t^2 + 1)$
6. $\mathbf{I}_2^* \mathbf{II}$ $y^2 = x^3 + t(t+2)x^2 + t^5(2t+1)$
 $-\Delta = t^8(t+2)^3(2t+1)$
7. $\mathbf{I}_2^* \mathbf{II} \mathbf{1}$ $y^2 = x^3 + t(t+2)x^2 + t^5(2t+2)$
 $-\Delta = t^8(t+2)^3(2t+2)$

4.2.16 Case 5C

$$\delta = 9, \quad r = 7 \quad \text{Type } \mathbf{I}_3^*$$

$$y^2 = x^3 + tc_1x^2 + t^4c_0x + t^6d_0, \quad t \nmid c_1, t \nmid d_0 \quad (4.2.16)$$

$$-\Delta = t^2c_1^2(t^7c_1d_0 - t^8c_0^2) + t^{12}c_0^3$$

1. $\mathbf{I}_3^* \mathbf{3}$ does not exist, A
2. $\mathbf{I}_3^* \mathbf{12}$ does not exist, C
3. $\mathbf{I}_3^* \mathbf{1^3}$ $y^2 = x^3 + tx^2 + 2t^4x + 2t^6$
 $-\Delta = t^{12} + t^{10} + t^9 = t^9(t-1)(t^2 + t + 2)$
4. $\mathbf{I}_3^* \mathbf{II}$ $y^2 = x^3 + t(t+2)x^2 + 2t^6$
 $-\Delta = 2t^9(t+2)^3$
5. $\mathbf{I}_3^* \mathbf{III}$ does not exist, C

4.2.17 Case 5D

$$\delta = 10, \quad r = 8 \quad \text{Type } \mathbf{I}_4^*$$

$$\begin{aligned} y^2 &= x^3 + tc_1x^2 + t^4c_0x \quad t \nmid c_0, t \nmid c_1 \\ -\Delta &= -t^{10}c_1^2c_0^2 + t^{12}c_0^3 \end{aligned} \quad (4.2.17)$$

1. $\mathbf{I}_4^* \mathbf{2}$ does not exist, A
2. $\mathbf{I}_4^* \mathbf{1^2}$ $y^2 = x^3 + tx^2 + t^4x$
 $-\Delta = t^{10}(t^2 - 1) = t^{10}(t+1)(t-1)$

4.2.18 Case 6A

$$\delta = 9, \quad r = 6 \quad \text{Type } \mathbf{IV}^*$$

$$\begin{aligned} y^2 &= x^3 + t^2c_0x^2 + t^3c_1x + t^4c_2, \quad t \nmid c_2, t \nmid c_1 \\ -\Delta &= t^4c_0^2(t^6c_0c_2 - t^6d_1^2) + t^9c_1^3 \end{aligned} \quad (4.2.18)$$

1. $\mathbf{IV}^* \mathbf{3}$ $y^2 = x^3 + t^2x^2 + t^3x + t^4(2t^2 + 1)$
 $-\Delta = t^{12} - t^9 = t^9(t-1)^3$
2. $\mathbf{IV}^* \mathbf{21}$ $y^2 = x^3 + t^2x^2 + 2t^3x + 2t^6 + t^5 + 2t^4$
 $-\Delta = t^{12} + 2t^{11} + 2t^{10} + t^9 = t^9(t-1)^2(t+1)$
3. $\mathbf{IV}^* \mathbf{1^3}$ $y^2 = x^3 + t^2x^2 + t^4x + 2t^6$
 $-\Delta = t^{12} + t^{11} + 2t^{10} + t^9 = t^9(t^3 + t^2 - t + 1)$
4. $\mathbf{IV}^* \mathbf{II}$ $y^2 = x^3 + t^3(2t+1)x + t^4$
 $-\Delta = t^9(2t+1)^3$
5. $\mathbf{IV}^* \mathbf{III}$ $y^2 = x^3 + t^3(t+2)x + t^4(t^2 + t + 1)$
 $-\Delta = t^9(t+2)^3$

4.2.19 Case 6B

$$\delta = 10, \quad r = 6 \quad \text{Type IV}^*$$

$$\begin{aligned} y^2 &= x^3 + t^2 c_0 x^2 + t^4 d_0 x + t^4 c_2 \quad t \nmid c_2, t \nmid c_0 \\ -\Delta &= t^4 c_0^2 (t^6 c_0 c_2 - t^8 d_0^2) + t^{12} d_0^3 \end{aligned} \quad (4.2.19)$$

1. **IV* 1²** $y^2 = x^3 + tx^2 + t^4(2t^2 + 1)$
 $-\Delta = t^{12} + 2t^{10} = t^{10}(t+1)(t-1)$
2. **IV* 2** $y^2 = x^3 + t^2 c_0 x^2 + t^4(\alpha t^2 + \alpha t + \alpha)$
 $-\Delta = t^{10}(c_0^3 \alpha (t-1)^2)$

4.2.20 Case 6C

$$\delta = 12, \quad r = 6 \quad \text{Type IV}^*$$

$$\begin{aligned} y^2 &= x^3 + t^4 d_0 x + t^4 c_2, \quad t \nmid c_2, t \nmid d_0 \\ -\Delta &= t^{12} d_0^3 \end{aligned} \quad (4.2.20)$$

4.2.21 Case 7

$$\delta = 9, \quad r = 7 \quad \text{Type III}^*$$

$$\begin{aligned} y^2 &= x^3 + t^2 c_0 x^2 + t^3 c_1 x + t^5 d_1, \quad t \nmid c_1 \\ -\Delta &= t^4 c_0^2 (t^7 c_0 d_1 - t^6 c_1^2) + t^9 c_1^3 \end{aligned} \quad (4.2.21)$$

1. **III* 3** does not exist, A
2. **III* 2 1** $y^2 = x^3 + t^2 x^2 + t^3(t+1)x + t^6 + t^5$
 $-\Delta = t^{12} + 2t^{11} + 2t^{10} + t^9 = t^9(t+1)(t^2 + t + 1)$
3. **III* 1³** $y^2 = x^3 + t^2 x^2 + t^3(t+1)x + 2t^6$
 $-\Delta = 2t^{12} + t^{11} + 2t^{10} + t^9 = t^9(2t+1)(t^2 + 1)$
4. **III* II** $y^2 = x^3 + t^3(2t+1)x + t^5(2+t)$
 $-\Delta = t^9(2t+1)^3$
5. **III* III** $y^2 = x^3 + t^3(2t+1)x$
 $-\Delta = t^9(2t+1)^3$

4.2.22 Case 8A

$$\delta = 11, \quad r = 8 \quad \text{Type II}^*$$

$$\begin{aligned} y^2 &= x^3 + t^2 c_0 x^2 + t^4 d_0 x + t^5 d_1, \quad t \nmid d_1, t \nmid c_0 \\ -\Delta &= t^4 c_0^2 (t^7 c_0 d_1 - t^8 d_0^2) + t^{12} d_0^3 \end{aligned} \quad (4.2.22)$$

1. $\text{II}_2^* 1 \quad y^2 = t^2 x^2 + t^4 x + t^5$
 $-\Delta = t^{11} (t + 1)$

4.2.23 Case 8B

$$\delta = 12, \quad r = 8 \quad \text{Type II}^*$$

$$\begin{aligned} y^2 &= x^3 + t^4 d_0 x + t^5 d_1, \quad t \nmid d_1, t \nmid d_0 \\ -\Delta &= t^{12} d_0^3 \end{aligned} \quad (4.2.23)$$

5 Conclusion and Summary of Results

5.1 General Results

Of the 372 configurations checked above, having either multiplicative or additive singularities, 267 configurations were found to exist in characteristic three, and 105 were found not to exist. 227 configurations involving at least one additive singularity exist, while 68 involving at least one additive singularity did not. Of those configurations which are purely multiplicative, 40 exist, while 37 do not.

5.2 Multiplicative Results in Comparison with other Characteristics

As mentioned earlier, in different characteristics the additive singularities have such different properties that their configurations are not easily compared from characteristic to characteristic. However, this is not a problem with purely multiplicative fibres, so we compare them here.

There are 6 configurations that exist in characteristic zero, but not in characteristic three. But every configuration that exists in characteristic three also exists in characteristic zero. There are 6 configurations which exist in characteristic two but not characteristic three, and there are 10 configurations that exist in characteristic three but not characteristic two.

The following table summarizes the results. A $\checkmark$ indicates that a rational elliptic surface with the indicated fibre distribution exists in the given case and characteristic, whereas an X indicates that no such surface exists.

Table 3: Existence of multiplicative fibre types

Partition	Characteristic 0	Characteristic 2	Characteristic 3
1^{12}	✓	✓	✓
21^{10}	✓	✓	✓
31^9	✓	✓	✓
$2^2 1^8$	✓	✓	✓
41^8	✓	✓	✓
321^7	✓	✓	✓
51^7	✓	✓	✓
$2^3 1^6$	✓	✓	✓
421^6	✓	✓	✓
$3^2 1^6$	✓	✓	✓
61^6	✓	✓	✓
$32^2 1^5$	✓	✓	✓
521^5	✓	✓	✓
431^5	✓	✓	✓
71^5	✓	✓	✓
$2^4 1^4$	✓	✓	✓
$42^2 1^4$	✓	✓	✓
$3^2 21^4$	✓	✓	✓
621^4	✓	✓	✓
531^4	✓	✓	✓
$4^2 1^4$	✓	✓	✓
81^4	✓	✓	✓
$32^3 1^3$	✓	✓	✓
$52^2 1^3$	✓	✓	✓
4321^3	✓	✓	✓
721^3	✓	✓	✓
$3^3 1^3$	✓	✓	X
631^3	✓	✓	X
541^3	✓	✓	✓
91^3	✓	✓	X
$2^5 1^2$	✓	X	✓
$42^3 1^2$	✓	X	✓
$3^2 2^2 1^2$	✓	✓	✓
$62^2 1^2$	✓	X	✓
5321^2	✓	✓	✓
$4^2 21^2$	✓	X	✓
821^2	✓	X	✓
$43^2 1^2$	X	X	X
731^2	X	X	X
641^2	X	X	X
<i>continued on next page</i>			

<i>continued from previous page</i>			
Partition	Characteristic 0	Characteristic 2	Characteristic 3
$5^2 1^2$	✓	✓	✓
$10 1^2$	X	X	X
$3 2^4 1$	✓	X	✓
$5 2^3 1$	X	X	X
$4 3 2^2 1$	✓	X	✓
$7 2^2 1$	X	X	X
$3^3 2 1$	✓	✓	X
$6 3 2 1$	✓	X	X
$5 4 2 1$	X	X	X
$9 2 1$	X	X	X
$5 3^2 1$	X	X	X
$4^2 3 1$	X	X	X
$8 3 1$	X	X	X
$7 4 1$	X	X	X
$6 5 1$	X	X	X
$11 1$	X	X	X
2^6	✓	X	✓
$4 2^4$	✓	X	✓
$3^2 2^3$	X	X	X
$6 2^3$	X	X	X
$5 3 2^2$	X	X	X
$4^2 2^2$	✓	X	✓
$8 2^2$	X	X	X
$4 3^2 2$	X	X	X
$7 3 2$	X	X	X
$6 4 2$	X	X	X
$5^2 2$	X	X	X
$10 2$	X	X	X
3^4	✓	✓	X
$6 3^2$	X	X	X
$5 4 3$	X	X	X
$9 3$	X	X	X
4^3	X	X	X
$8 4$	X	X	X
$7 5$	X	X	X
6^2	X	X	X
12	X	X	X

References

- [1] E. B. Dynkin, *Semisimple subalgebras of semisimple Lie algebras*, Mat. Sbornik N.S., **Vol. 30(72)** 1952, 349–462.
- [2] W. E. Lang, *Configurations of singular fibres on rational elliptic surfaces in characteristic two*, Communications in Algebra, **Vol. 28(12)** 2000, 5813–5836.
- [3] W. E. Lang, *Extremal rational elliptic surfaces in characteristic p. II: Surfaces with three or fewer singular fibres*, Arkiv for matematik, **Vol. 32** 1994, 423–438.
- [4] R. Miranda, *Persson’s list of singular fibers for a rational elliptic surface*, Mathematische Zeitschrift, **Vol. 205** 1990, 191–211.
- [5] R. Miranda and U. Persson, *On extremal rational elliptic surfaces*, Mathematische Zeitschrift, **Vol. 193** 1986, 537–558.
- [6] K. Oguiso and T. Shioda, *The Mordell-Weil lattice of a rational elliptic surface*, Comment. Math. Univ. St. Paul, **Vol. 40(1)** 1990, 83–99.
- [7] U. Persson, *Configurations of Kodaira fibers on rational elliptic surfaces*, Mathematische Zeitschrift, **Vol. 205(1)** 1990, 1–47.
- [8] T. Shioda, *On the Mordell-Weil lattices*, Comment. Math. Univ. St. Paul, **Vol. 39(2)** 1990, 211–240.
- [9] J. H. Silverman, *Advanced topics in the arithmetic of elliptic curves*, Graduate Texts in Mathematics, **Vol. 151** 1994.